

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|   |  |  |
|---|--|--|
| <p>(51) International Patent Classification <sup>7</sup> :<br/>A61K 31/165, C07C 239/18, A61K 31/395, 31/335, A61P 31/04, C07D 295/10, 233/64, 213/50, 307/81, 317/40, 307/52, 207/32, C07C 311/08, C07D 239/42, 239/12</p>   | <p>A1</p>  | <p>(11) International Publication Number: WO 00/61134</p>              |
|   |  | <p>(43) International Publication Date: 19 October 2000 (19.10.00)</p> |
| <p>(21) International Application Number: PCT/GB00/01337</p>  | <p>maceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB). BECKETT, Raymond, Paul [GB/GB]; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB).</p>                     |  |
| <p>(22) International Filing Date: 10 April 2000 (10.04.00)</p>   |  |  |
| <p>(30) Priority Data:<br/>9907981.6 9 April 1999 (09.04.99) GB<br/>9918984.7 11 August 1999 (11.08.99) GB<br/>9927089.4 16 November 1999 (16.11.99) GB</p>   | <p>(74) Agent: WALLS, Alan, J.; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB).</p>   |  |
| <p>(71) Applicant (for all designated States except US): BRITISH BIOTECH PHARMACEUTICALS LIMITED [GB/GB]; Watlington Road, Cowley, Oxford OX4 5LY (GB).</p>   | <p>(81) Designated States: AU, BR, CA, CN, CZ, DE, GB, GE, HU, IL, JP, KR, MX, NO, NZ, PL, RU, SG, SK, TR, UA, US, ZA, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> |  |
| <p>(72) Inventors; and<br/>(75) Inventors/Applicants (for US only): TODD, Richard, Simon [GB/GB]; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB). BROOKINGS, Daniel, Christopher [GB/GB]; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB). SMITH, Helen, Katherine [GB/GB]; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB). THOMPSON, Alison, Jane [GB/GB]; British Biotech Phar-</p> | <p>Published<br/>With international search report.</p>   |  |
| <p>(54) Title: ANTIMICROBIAL AGENTS</p>   |  |  |
| <div style="text-align: center;"> <p>(I)</p> </div>   |  |  |
| <p>(57) Abstract</p>  |  |  |
| <p>Compounds of formula (I) are antibacterial agents wherein: Z represents a radical of formula -N(OH)CH(=O) of formula -C(=O)NH(OH) and R<sub>1</sub>-R<sub>4</sub> are as defined in the specification.</p>   |  |  |

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                          |    |  |    |  |    |                          |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania                  | ES | Spain                                    | LS | Lesotho                                      | SI | Slovenia                 |
| AM | Armenia                  | FI | Finland                                  | LT | Lithuania                                    | SK | Slovakia                 |
| AT | Austria                  | FR | France                                   | LU | Luxembourg                                   | SN | Senegal                  |
| AU | Australia                | GA | Gabon                                    | LV | Latvia                                       | SZ | Swaziland                |
| AZ | Azerbaijan               | GB | United Kingdom                           | MC | Monaco                                       | TD | Chad                     |
| BA | Bosnia and Herzegovina   | GE | Georgia                                  | MD | Republic of Moldova                          | TG | Togo                     |
| BB | Barbados                 | GH | Ghana                                    | MG | Madagascar                                   | TJ | Tajikistan               |
| BE | Belgium                  | GN | Guinea                                   | MK | The former Yugoslav<br>Republic of Macedonia | TM | Turkmenistan             |
| BF | Burkina Faso             | GR | Greece                                   | ML | Mali   | TR | Turkey                   |
| BG | Bulgaria                 | HU | Hungary                                  | MN | Mongolia                                     | TT | Trinidad and Tobago      |
| BJ | Benin                    | IE | Ireland                                  | MR | Mauritania                                   | UA | Ukraine                  |
| BR | Brazil                   | IL | Israel                                   | MW | Malawi                                       | UG | Uganda                   |
| BY | Belarus                  | IS | Iceland                                  | MX | Mexico                                       | US | United States of America |
| CA | Canada                   | IT | Italy                                    | NE | Niger  | UZ | Uzbekistan               |
| CF | Central African Republic | JP | Japan                                    | NL | Netherlands                                  | VN | Viet Nam                 |
| CG | Congo                    | KE | Kenya                                    | NO | Norway                                       | YU | Yugoslavia               |
| CH | Switzerland              | KG | Kyrgyzstan                               | NZ | New Zealand                                  | ZW | Zimbabwe                 |
| CI | Côte d'Ivoire            | KP | Democratic People's<br>Republic of Korea | PL | Poland                                       |    |                          |
| CM | Cameroon                 | KR | Republic of Korea                        | PT | Portugal                                     |    |                          |
| CN | China                    | KZ | Kazakhstan                               | RO | Romania                                      |    |                          |
| CU | Cuba                     | LC | Saint Lucia                              | RU | Russian Federation                           |    |                          |
| CZ | Czech Republic           | LJ | Liechtenstein                            | SD | Sudan  |    |                          |
| DE | Germany                  | LK | Sri Lanka                                | SE | Sweden                                       |    |                          |
| DK | Denmark                  | LR | Liberia                                  | SG | Singapore                                    |    |                          |
| EE | Estonia                  |    |  |    |  |    |                          |

## Antimicrobial Agents

This invention relates to a novel class of hydroxamic acid and N-formyl hydroxylamine derivatives having antibacterial activity, and to pharmaceutical and veterinary compositions comprising such compounds.

### Background to the Invention

In general, bacterial pathogens are classified as either Gram-positive or Gram-negative. Many antibacterial agents (including antibiotics) are specific against one or other Gram-class of pathogens. Antibacterial agents effective against both Gram-positive and Gram-negative pathogens are therefore generally regarded as having broad spectrum activity.

Many classes of antibacterial agents are known, including the penicillins and cephalosporins, tetracyclines, sulfonamides, monobactams, fluoroquinolones and quinolones, aminoglycosides, glycopeptides, macrolides, polymyxins, lincosamides, trimethoprim and chloramphenicol. The fundamental mechanisms of action of these antibacterial classes vary.

Bacterial resistance to many known antibacterials is a growing problem. Accordingly there is a continuing need in the art for alternative antibacterial agents, especially those which have mechanisms of action fundamentally different from the known classes.

Amongst the Gram-positive pathogens, such as Staphylococci, Streptococci, Mycobacteria and Enterococci, resistant strains have evolved/arisen which makes them particularly difficult to eradicate. Examples of such strains are methicillin resistant *Staphylococcus aureus* (MRSA), methicillin resistant coagulase negative Staphylococci (MRCNS), penicillin resistant *Streptococcus pneumoniae* and multiply resistant *Enterococcus faecium*.

Pathogenic bacteria are often resistant to the aminoglycoside,  $\beta$ -lactam (penicillins and cephalosporins), and chloramphenicol types of antibiotic. This resistance involves the enzymatic inactivation of the antibiotic by hydrolysis or by formation of inactive derivatives. The  $\beta$ -lactam (penicillin and cephalosporin) family of antibiotics are characterised by the presence of a  $\beta$ -lactam ring structure. Resistance to this family of antibiotics in clinical isolates is most commonly due to the production of a "penicillinase" ( $\beta$ -lactamase) enzyme by the resistant bacterium which hydrolyses the  $\beta$ -lactam ring thus eliminating its antibacterial activity.

Recently there has been an emergence of vancomycin-resistant strains of enterococci (Woodford N. 1998 Glycopeptide-resistant enterococci: a decade of experience. Journal of Medical Microbiology. 47(10):849-62). Vancomycin-resistant enterococci are particularly hazardous in that they are frequent causes of hospital based infections and are inherently resistant to most antibiotics. Vancomycin works by binding to the terminal D-Ala-D-Ala residues of the cell wall peptidoglycan precursor. The high-level resistance to vancomycin is known as VanA and is conferred by a genes located on a transposable element which alter the terminal residues to D-Ala-D-lac thus reducing the affinity for vancomycin.

In view of the rapid emergence of multidrug-resistant bacteria, the development of antibacterial agents with novel modes of action that are effective against the growing number of resistant bacteria, particularly the vancomycin resistant enterococci and  $\beta$ -lactam antibiotic-resistant bacteria, such as methicillin-resistant *Staphylococcus aureus*, is of utmost importance.

#### Brief Description of the Invention

This invention is based on the finding that certain hydroxamic acid and N-formyl hydroxylamine derivatives have antibacterial activity, and makes available a new class of antibacterial agents. The inventors have found that the compounds with which this invention is concerned are antibacterial with respect to a range of Gram-positive and Gram-negative organisms.



Although it may be of interest to establish the mechanism of action of the compounds with which the invention is concerned, it is their ability to inhibit bacterial growth that makes them useful. However, it is presently believed that their antibacterial activity is due, at least in part, to intracellular inhibition of bacterial polypeptide deformylase (PDF; EC 3.5.1.31).

All ribosome-mediated synthesis of proteins starts with a methionine residue. In prokaryotes the methionyl moiety carried by the initiator tRNA is N-formylated prior to its incorporation into a polypeptide. Consequently, N-formylmethionine is always present at the N-terminus of a nascent bacterial polypeptide. However, most mature proteins do not retain the N-formyl group or the terminal methionine residue. Deformylation is required prior to methionine removal, since methionine aminopeptidase does not recognise peptides with an N-terminal formylmethionine residue (Solbiati et al., J. Mol. Biol. 290:607-614, 1999). Deformylation is, therefore, a crucial step in bacterial protein biosynthesis and the enzyme responsible, PDF, is essential for normal bacterial growth. Although the gene encoding PDF (*def*) is present in all pathogenic bacteria for which sequences are known (Meinzel et al., J. Mol. Biol., 266:939-49, 1997), it has no eukaryotic counterpart, making it an attractive target for antibacterial chemotherapy.

The isolation and characterisation of PDF has been facilitated by an understanding of the importance of the metal ion in the active site (Groche et al., Biophys. Biochem. Res. Commun., 246:324-6, 1998). The Fe<sup>2+</sup> form is highly active *in vivo* but is unstable when isolated due to oxidative degradation (Rajagopalan et al., J. Biol. Chem. 273:22305-10, 1998). The Ni<sup>2+</sup> form of the enzyme has specific activity comparable with the ferrous enzyme but is oxygen-insensitive (Ragusa et al., J. Mol. Biol. 1998, 280:515-23, 1998). The Zn<sup>2+</sup> enzyme is also stable but is almost devoid of catalytic activity (Rajagopalan et al., J. Am. Chem. Soc. 119:12418-12419, 1997).

Several X-ray crystal structures and NMR structures of *E. coli* PDF, with or without

bound inhibitors, have been published (Chan et al., Biochemistry 36:13904-9, 1997; Becker et al., Nature Struct. Biol. 5:1053-8, 1998; Becker et al., J. Biol. Chem. 273:11413-6, 1998; Hao et al., Biochemistry, 38:4712-9, 1999; Dardel et al., J. Mol. Biol. 280:501-13, 1998; O'Connell et al., J. Biomol. NMR, 13:311-24, 1999), indicating similarities in active site geometry to metalloproteinases such as thermolysin and the metzincins.

Recently the substrate specificity of PDF has been extensively studied (Ragusa et al., J. Mol. Biol. 289:1445-57, 1999; Hu et al., Biochemistry 38:643-50, 1999; Meinnel et al., Biochemistry, 38:4287-95, 1999). These authors conclude that an unbranched hydrophobic chain is preferred at P1', while a wide variety of P2' substituents are acceptable and an aromatic substituent may be advantageous at the P3' position. There have also been reports that small peptidic compounds containing an H-phosphonate (Hu et al., Bioorg. Med. Chem. Lett., 8:2479-82, 1998) or thiol (Meinnel et al., Biochemistry, 38:4287-95, 1999) metal binding group are micromolar inhibitors of PDF. Peptide aldehydes such as calpeptin (N-Cbz-Leu-norleucinal) have also been shown to inhibit PDF (Durand et al., Arch. Biochem. Biophys., 367:297-302, 1999). However, the identity of the metal binding group and its spacing from the rest of the molecule ("recognition fragment") has not been studied extensively. Furthermore, non-peptidic PDF inhibitors, which may be desirable from the point of view of bacterial cell wall permeability or oral bioavailability in the host species, have not been identified.

Recently it has been reported that PDF is present in eukaryotic parasites such as *Plasmodium falciparum* (Ferreira et al, Parasitology Today, vol 16, no. 4, 2000). Those authors also found evidence for the presence of PDF in other parasites of humans, such as the kinetoplastid protozoan parasites *Trypanosoma brucei* and *Leishmania major*. Based on these findings, it is anticipated that the compounds with which this invention is concerned have antiprotozoal activity, and are useful in the treatment of malaria and other protozoal diseases.


Certain N-formyl hydroxylamine derivatives have previously been claimed in the patent publications listed below, although very few examples of such compounds have been specifically made and described:

|              |                       |
|--------------|-----------------------|
| EP-B-0236872 | (Roche)               |
| WO 92/09563  | (Glycomed)            |
| WO 92/04735  | (Syntex)              |
| WO 95/19965  | (Glycomed)            |
| WO 95/22966  | (Sanofi Winthrop)     |
| WO 95/33709  | (Roche)               |
| WO 96/23791  | (Syntex)              |
| WO 96/16027  | (Syntex/Agouron)      |
| WO 97/03783  | (British Biotech)     |
| WO 97/18207  | (DuPont Merck)        |
| WO 98/38179  | (GlaxoWellcome)       |
| WO 98/47863  | (Labs Jaques Logeais) |

The pharmaceutical utility ascribed to the N-formyl hydroxylamine derivatives in those publications is the ability to inhibit matrix metalloproteinases (MMPs) and in some cases release of tumour necrosis factor (TNF), and hence the treatment of diseases or conditions mediated by those enzymes, such as cancer and rheumatoid arthritis. That prior art does not disclose or imply that N-formyl hydroxylamine derivatives have antibacterial activity.

In addition to these, US-A-4,738,803 (Roques et al.) also discloses N-formyl hydroxylamine derivatives, however, these compounds are disclosed as enkephalinase inhibitors and are proposed for use as antidepressants and hypotensive agents. Also, WO 97/38705 (Bristol-Myers Squibb) discloses certain N-formyl hydroxylamine derivatives as enkephalinase and angiotensin converting enzyme inhibitors. This prior art does not disclose or imply that N-formyl hydroxylamine derivatives have antibacterial activity either.

Our copending International patent applications nos. WO 99/39704 and WO 99/59568 disclose that certain N-formyl hydroxylamine and hydroxamic acid derivatives have antibacterial activity. One class of compounds disclosed therein as having such activity has general formula (IA):




Wherein X represents an N-formylhydroxylamino group (-N(OH)CHO) or a hydroxamic acid group (-CONHOH), and the various "R" substituents are as specified in the documents. The compounds useful in accordance with the present invention differ in structure from those of WO 99/39704 and WO 99/59568 principally in that the amido radical  $R_6R_5NCO-$  is replaced by a ketone radical.

Very many hydroxamic acid derivatives are known. Many have been disclosed as having matrix metalloproteinase (MMP) inhibitory activity, and thus to be potentially useful for the treatment of diseases mediated by MMPs, for example cancer, arthritides, and conditions involving tissue remodeling such as wound healing, and restenosis. However, very few such known hydroxamic acid derivatives have the ketone moiety which characterises the compounds useful according to the present invention. One publication which does, however, is WO 98/30541 (Abbott).

#### Detailed description of the invention

According to the first aspect of the present invention there is provided the use of a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt, hydrate or solvate thereof in the preparation of a composition for treatment of

bacterial or protozoal infections in humans and non-human mammals:



wherein:

Z represents a radical of formula  $-N(OH)CH(=O)$  or formula  $-C(=O)NH(OH)$ ;

$R_1$  represents hydrogen, methyl or trifluoromethyl, or, except when Z is a radical of formula  $-N(OH)CH(=O)$ , a hydroxy or amino group;

$R_2$  represents a group  $R_{10}-(X)_n-(ALK)_m$ - wherein

$R_{10}$  represents hydrogen, or a  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl, cycloalkyl, aryl, or heterocyclyl group, any of which may be unsubstituted or substituted by  $(C_1$ - $C_6)$ alkyl,  $(C_1$ - $C_6)$ alkoxy, hydroxy, mercapto,  $(C_1$ - $C_6)$ alkylthio, amino, halo (including fluoro, chloro, bromo and iodo), trifluoromethyl, cyano, nitro, oxo,  $-COOH$ ,  $-CONH_2$ ,  $-COOR^A$ ,  $-NHCOR^A$ ,  $-CONHR^A$ ,  $-NHR^A$ ,  $-NR^A R^B$ , or  $-CONR^A R^B$  wherein  $R^A$  and  $R^B$  are independently a  $(C_1$ - $C_6)$ alkyl group and

ALK represents a straight or branched divalent  $C_1$ - $C_6$  alkylene,  $C_2$ - $C_6$  alkenylene, or  $C_2$ - $C_6$  alkynylene radical, and may be interrupted by one or more non-adjacent  $-NH-$ ,  $-O-$  or  $-S-$  linkages,

X represents  $-NH-$ ,  $-O-$  or  $-S-$ , and

m and n are independently 0 or 1;

$R_3$  represents the side chain of a natural or non-natural alpha amino acid; and

$R_4$  represents a radical  $R_5-(ALK)_p$ - wherein ALK is as defined in relation to  $R_2$ , p is 0 or 1, and  $R_5$  represents hydrogen, or a  $C_1-C_6$  alkyl,  $C_2-C_6$  alkenyl,  $C_2-C_6$  alkynyl, cycloalkyl, aryl, or heterocyclyl group any of which

(i) may be substituted by a group selected from ( $C_1-C_6$ )alkyl, phenyl, benzyl, ( $C_1-C_6$ )alkoxy, phenoxy, hydroxy, mercapto, ( $C_1-C_6$ )alkylthio, amino, halo, trifluoromethyl, cyano, nitro, oxo,  $-COOH$ ,  $-SO_2H$ ,  $-CONH_2$ ,  $-SO_2NH_2$ ,  $-COR^A$ ,  $-SOR^A$ ,  $-SO_2R^A$ ,  $-COOR^A$ ,  $-CONHR^A$ ,  $-SO_2NHR^A$ ,  $-NHCOR^A$ ,  $-NHSO_2R^A$ , and  $-NHR^A$ , wherein  $R^A$  is

( $C_1-C_6$ )alkyl, cycloalkyl, phenyl, 2-, 3- or 4-pyridyl, N- or 2-, 3- or 4-piperidyl, N- or 2- or 3- piperazyl group; or

(ii) may be substituted by  $-NR^AR^B$ ,  $-CONR^AR^B$  or  $-SO_2NR^AR^B$ , wherein  $R^A$  and  $R^B$  are independently

( $C_1-C_6$ )alkyl, cycloalkyl, phenyl, 2-, 3- or 4-pyridyl, N- or 2-, 3- or 4-piperidyl, N- or 2- or 3- piperazyl group,

or when taken together with the N atom to which they are attached  $R^A$  and  $R^B$  form a 5 to 7 membered aromatic or non- aromatic ring, which ring (a) may contain additional heteroatoms selected from N, O and S, and in which any S atom may be oxidised as a sulphonyl or sulphoxide, and (b) may be substituted on a ring carbon or heteroatom by one or more of the substituents listed under (i) above.

In another aspect, the invention provides a method for the treatment of bacterial or protozoal infections in humans and non-human mammals, which comprises administering to a subject suffering such infection an antibacterially effective dose of a compound of formula (I) as defined above.

In a further aspect of the invention there is provided a method for the treatment of

bacterial contamination by applying an antibacterially effective amount of a compound of formula (I) as defined above to the site of contamination.

The compounds of formula (I) as defined above may be used as component(s) of antibacterial cleaning or disinfecting materials.

Compounds of formula (I) above wherein Z is a radical of formula  $-N(OH)CH(=O)$ , are believed to be novel. Accordingly, in another aspect, the invention provides a compound of formula (I) wherein Z is a radical of formula  $-N(OH)CH(=O)$ , or a pharmaceutically or veterinarily acceptable salt, hydrate or solvate thereof.

On the hypothesis that the compounds (I) act by inhibition of intracellular PDF, the most potent antibacterial effect may be achieved by using compounds which efficiently pass through the bacterial cell wall. Thus, compounds which are highly active as inhibitors of PDF in vitro and which penetrate bacterial cells are preferred for use in accordance with the invention. It is to be expected that the antibacterial potency of compounds which are potent inhibitors of the PDF enzyme in vitro, but are poorly cell penetrant, may be improved by their use in the form of a prodrug, ie a structurally modified analogue which is converted to the parent molecule of formula (I), for example by enzymic action, after it has passed through the bacterial cell wall. The same is true in the case of protozoa.

As used herein the term " $(C_1-C_6)$ alkyl" means a straight or branched chain alkyl moiety having from 1 to 6 carbon atoms, including for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl and n-hexyl.

As used herein the term "divalent  $(C_1-C_6)$ alkylene radical" means a saturated hydrocarbon chain having from 1 to 6 carbon atoms and two unsatisfied valencies.

As used herein the term " $(C_2-C_6)$ alkenyl" means a straight or branched chain alkenyl moiety having from 2 to 6 carbon atoms having at least one double bond of either E

or Z stereochemistry where applicable. The term includes, for example, vinyl, allyl, 1- and 2-butenyl and 2-methyl-2-propenyl.

As used herein the term "divalent ( $C_2-C_6$ )alkenylene radical" means a hydrocarbon chain having from 2 to 6 carbon atoms, at least one double bond, and two unsatisfied valencies.

As used herein the term " $C_2-C_6$  alkynyl" refers to straight chain or branched chain hydrocarbon groups having from two to six carbon atoms and having in addition one triple bond. This term would include for example, ethynyl, 1-propynyl, 1- and 2-butenyl, 2-methyl-2-propynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl and 5-hexynyl.

As used herein the term "divalent ( $C_2-C_6$ )alkynylene radical" means a hydrocarbon chain having from 2 to 6 carbon atoms, at least one triple bond, and two unsatisfied valencies.

As used herein the term "cycloalkyl" means a saturated alicyclic moiety having from 3-8 carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

As used herein the term "cycloalkenyl" means an unsaturated alicyclic moiety having from 3-8 carbon atoms and includes, for example, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl and cyclooctenyl. In the case of cycloalkenyl rings of from 5-8 carbon atoms, the ring may contain more than one double bond.

As used herein the term "aryl" refers to a mono-, bi- or tri-cyclic carbocyclic aromatic group, and to groups consisting of two covalently linked monocyclic carbocyclic aromatic groups. Illustrative of such groups are phenyl, biphenyl and naphthyl.



As used herein the unqualified term "heterocyclyl" or "heterocyclic" includes "heteroaryl" as defined below, and in particular means a 5-8 membered aromatic or non-aromatic heterocyclic ring containing one or more heteroatoms selected from S, N and O, and optionally fused to a benzyl or second heterocyclic ring, and the term includes, for example, pyrrolyl, furyl, thienyl, piperidinyl, imidazolyl, oxazolyl, thiazolyl, thiadiazolyl, thiazepinyl, pyrazolyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl, and benzimidazolyl rings.

As used herein the term "heteroaryl" refers to a 5- or 6- membered aromatic ring containing one or more heteroatoms, and optionally fused to a benzyl or pyridyl ring; and to groups consisting of two covalently linked 5- or 6- membered aromatic rings each containing one or more heteroatoms; and to groups consisting of a monocyclic carbocyclic aromatic group covalently linked to a 5- or 6- membered aromatic rings containing one or more heteroatoms. Illustrative of such groups are thienyl, furyl, pyrrolyl, imidazolyl, benzimidazolyl, thiazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, 4-([1,2,3]-thiadiazol-4-yl)phenyl and 5-isoxazol-3-ylthienyl.

As used herein the unqualified term "carbocyclyl" or "carbocyclic" refers to a 5-8 membered ring whose ring atoms are all carbon.

Unless otherwise specified in the context in which it occurs, the term "substituted" as applied to any moiety herein means substituted with up to four substituents, each of which independently may be (C<sub>1</sub>-C<sub>8</sub>)alkyl, phenyl, benzyl, (C<sub>1</sub>-C<sub>8</sub>)alkoxy, phenoxy, hydroxy, mercapto, (C<sub>1</sub>-C<sub>8</sub>)alkylthio, amino, halo (including fluoro, chloro, bromo and iodo), trifluoromethyl, cyano, nitro, oxo, -COOH, -CONH<sub>2</sub>, -COR<sup>A</sup>, -COOR<sup>A</sup>, -NHCOR<sup>A</sup>, -CONHR<sup>A</sup>, -NHR<sup>A</sup>, -NR<sup>A</sup>R<sup>B</sup>, or -CONR<sup>A</sup>R<sup>B</sup> wherein R<sup>A</sup> and R<sup>B</sup> are independently a (C<sub>1</sub>-C<sub>8</sub>)alkyl group. In the case where "substituted" means substituted by benzyl, the phenyl ring thereof may itself be substituted with any of the foregoing, except phenyl or benzyl.

As used herein the terms "side chain of a natural alpha-amino acid" and "side chain of a non-natural alpha-amino acid" mean the group  $R^*$  in respectively a natural and non-natural amino acid of formula  $NH_2-CH(R^*)-COOH$ .

Examples of side chains of natural alpha amino acids include those of alanine, arginine, asparagine, aspartic acid, cysteine, cystine, glutamic acid, histidine, 5-hydroxylysine, 4-hydroxyproline, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine,  $\alpha$ -aminoadipic acid,  $\alpha$ -amino-n-butyric acid, 3,4-dihydroxyphenylalanine, homoserine,  $\alpha$ -methylserine, ornithine, pipecolic acid, and thyroxine.

In natural alpha-amino acid side chains which contain functional substituents, for example amino, carboxyl, hydroxy, mercapto, guanidyl, imidazolyl, or indolyl groups as in arginine, lysine, glutamic acid, aspartic acid, tryptophan, histidine, serine, threonine, tyrosine, and cysteine, such functional substituents may optionally be protected.

Likewise, in the side chains of non-natural alpha amino acids which contain functional substituents, for example amino, carboxyl, hydroxy, mercapto, guanidyl, imidazolyl, or indolyl groups, such functional substituents may optionally be protected.

The term "protected" when used in relation to a functional substituent in a side chain of a natural or non-natural alpha-amino acid means a derivative of such a substituent which is substantially non-functional. The widely used handbook by T. W. Greene and P. G. Wuts "Protective Groups in Organic Synthesis" Second Edition, Wiley, New York, 1991 reviews the subject. For example, carboxyl groups may be esterified (for example as a  $C_1-C_6$  alkyl ester), amino groups may be converted to amides (for example as a  $NHCOC_1-C_6$  alkyl amide) or carbamates (for example as an  $NHC(=O)OC_1-C_6$  alkyl or  $NHC(=O)OCH_2Ph$  carbamate), hydroxyl groups may be converted to ethers (for example an  $OC_1-C_6$  alkyl or a  $O(C_1-C_6$  alkyl)phenyl ether) or

esters (for example a  $\text{OC}(=\text{O})\text{C}_1\text{-C}_6$  alkyl ester) and thiol groups may be converted to thioethers (for example a tert-butyl or benzyl thioether) or thioesters (for example a  $\text{SC}(=\text{O})\text{C}_1\text{-C}_6$  alkyl thioester).

There are at least two actual or potential chiral centres in the compounds according to the invention because of the presence of asymmetric carbon atoms. The presence of several asymmetric carbon atoms gives rise to a number of diastereoisomers with R or S stereochemistry at each chiral centre. The invention includes all such diastereoisomers and mixtures thereof. Currently, the preferred stereoconfiguration of the carbon atom carrying the  $\text{R}_2$  group is R; that of the carbon atom carrying the  $\text{R}_1$  group (when asymmetric) is R; and that of the carbon atom carrying the  $\text{R}_3$  group (when asymmetric) is S

In the compounds of formula (I) as defined above:

$\text{R}_1$  is hydrogen, methyl, or trifluoromethyl. Hydrogen is currently preferred.

$\text{R}_2$  may be, for example:

optionally substituted  $\text{C}_1\text{-C}_6$  alkyl,  $\text{C}_3\text{-C}_6$  alkenyl,  $\text{C}_3\text{-C}_6$  alkynyl or cycloalkyl;

phenyl( $\text{C}_1\text{-C}_6$  alkyl)-, phenyl( $\text{C}_3\text{-C}_6$  alkenyl)- or phenyl( $\text{C}_3\text{-C}_6$  alkynyl)-  
optionally substituted in the phenyl ring;

cycloalkyl( $\text{C}_1\text{-C}_6$  alkyl)-, cycloalkyl( $\text{C}_3\text{-C}_6$  alkenyl)- or cycloalkyl( $\text{C}_3\text{-C}_6$  alkynyl)-  
optionally substituted in the cycloalkyl ring;

heterocyclyl( $\text{C}_1\text{-C}_6$  alkyl)-, heterocyclyl( $\text{C}_3\text{-C}_6$  alkenyl)- or heterocyclyl( $\text{C}_3\text{-C}_6$  alkynyl)- optionally substituted in the heterocyclyl ring; or

$\text{CH}_3(\text{CH}_2)_p\text{O}(\text{CH}_2)_q\text{-}$  or  $\text{CH}_3(\text{CH}_2)_p\text{S}(\text{CH}_2)_q\text{-}$ , wherein p is 0, 1, 2 or 3 and q is 1, 2 or 3.

Specific examples of  $R_2$  groups include

methyl, ethyl, n- and iso-propyl, n- and iso-butyl, n-pentyl, iso-pentyl 3-methyl-but-1-yl, n-hexyl, n-heptyl, n-octyl, methylsulfanylethyl, ethylsulfanylmethyl, 2-methoxyethyl, 2-ethoxyethyl, 2-ethoxymethyl, 3-hydroxypropyl, allyl, 3-phenylprop-3-en-1-yl, prop-2-yn-1-yl, 3-phenylprop-2-yn-1-yl, 3-(2-chlorophenyl)prop-2-yn-1-yl, but-2-yn-1-yl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclopentylethyl, cyclopentylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexylpropyl, furan-2-ylmethyl, furan-3-methyl, tetrahydrofuran-2-ylmethyl, tetrahydrofuran-2-ylmethyl, piperidinylmethyl, phenylpropyl, 4-chlorophenylpropyl, 4-methylphenylpropyl, 4-methoxyphenylpropyl, benzyl, 4-chlorobenzyl, 4-methylbenzyl, and 4-methoxybenzyl.

Presently preferred groups at  $R_2$  are n-propyl, n-butyl, n-pentyl, and cyclopentylmethyl.

$R_3$  may be, for example

the characterising group of a natural  $\alpha$  amino acid, for example benzyl, or 4-methoxyphenylmethyl, in which any functional group may be protected, any amino group may be acylated and any carboxyl group present may be amidated; or

a group  $-\text{[Alk]}_n\text{R}_9$  where Alk is a  $(\text{C}_1\text{-C}_6)$ alkylene or  $(\text{C}_2\text{-C}_6)$ alkenylene group optionally interrupted by one or more -O-, or -S- atoms or  $-\text{N}(\text{R}_{12})$ - groups [where  $\text{R}_{12}$  is a hydrogen atom or a  $(\text{C}_1\text{-C}_6)$ alkyl group], n is 0 or 1, and  $\text{R}_9$  is hydrogen or an optionally substituted phenyl, aryl, heterocyclyl, cycloalkyl or cycloalkenyl group or (only when n is 1)  $\text{R}_9$  may additionally be hydroxy, mercapto,  $(\text{C}_1\text{-C}_6)$ alkylthio, amino, halo, trifluoromethyl, nitro,  $-\text{COOH}$ ,  $-\text{CONH}_2$ ,  $-\text{COOR}^A$ ,  $-\text{NHCOR}^A$ ,  $-\text{CONHR}^A$ ,  $-\text{NHR}^A$ ,  $-\text{NR}^A\text{R}^B$ , or  $-\text{CONR}^A\text{R}^B$  wherein  $\text{R}^A$  and  $\text{R}^B$  are independently a  $(\text{C}_1\text{-C}_6)$ alkyl group; or

$\text{CONH}_2$ ,  $-\text{COOR}^A$ ,  $-\text{NHCOR}^A$ ,  $-\text{CONHR}^A$ ,  $-\text{NHR}^A$ ,  $-\text{NR}^A\text{R}^B$ , or  $-\text{CONR}^A\text{R}^B$

wherein  $\text{R}^A$  and  $\text{R}^B$  are independently a  $(\text{C}_1\text{-C}_6)$ alkyl group; or

a benzyl group substituted in the phenyl ring by a group of formula -  
 $\text{OCH}_2\text{COR}_8$ , where  $\text{R}_8$  is hydroxyl, amino,  $(\text{C}_1\text{-C}_6)$ alkoxy, phenyl $(\text{C}_1\text{-C}_6)$ alkoxy,  
 $(\text{C}_1\text{-C}_6)$ alkylamino, di $(\text{C}_1\text{-C}_6)$ alkylamino, phenyl $(\text{C}_1\text{-C}_6)$ alkylamino; or

a heterocyclic $(\text{C}_1\text{-C}_6)$ alkyl group, either being unsubstituted or mono- or di-  
substituted in the heterocyclic ring with halo, nitro, carboxy,  $(\text{C}_1\text{-C}_6)$ alkoxy,  
cyano,  $(\text{C}_1\text{-C}_6)$ alkanoyl, trifluoromethyl  $(\text{C}_1\text{-C}_6)$ alkyl, hydroxy, formyl, amino,  
 $(\text{C}_1\text{-C}_6)$ alkylamino, di $(\text{C}_1\text{-C}_6)$ alkylamino, mercapto,  $(\text{C}_1\text{-C}_6)$ alkylthio,  
hydroxy $(\text{C}_1\text{-C}_6)$ alkyl, mercapto $(\text{C}_1\text{-C}_6)$ alkyl or  $(\text{C}_1\text{-C}_6)$ alkylphenylmethyl; or

a group  $-\text{CR}_a\text{R}_b\text{R}_c$  in which:

each of  $\text{R}_a$ ,  $\text{R}_b$  and  $\text{R}_c$  is independently hydrogen,  $(\text{C}_1\text{-C}_6)$ alkyl,  $(\text{C}_2\text{-C}_6)$ alkenyl,  $(\text{C}_2\text{-C}_6)$ alkynyl, phenyl $(\text{C}_1\text{-C}_6)$ alkyl,  $(\text{C}_3\text{-C}_8)$ cycloalkyl; or

$\text{R}_c$  is hydrogen and  $\text{R}_a$  and  $\text{R}_b$  are independently phenyl or heteroaryl  
such as pyridyl; or

$\text{R}_c$  is hydrogen,  $(\text{C}_1\text{-C}_6)$ alkyl,  $(\text{C}_2\text{-C}_6)$ alkenyl,  $(\text{C}_2\text{-C}_6)$ alkynyl, phenyl $(\text{C}_1\text{-C}_6)$ alkyl, or  $(\text{C}_3\text{-C}_8)$ cycloalkyl, and  $\text{R}_a$  and  $\text{R}_b$  together with the carbon  
atom to which they are attached form a 3 to 8 membered cycloalkyl or  
a 5- to 6-membered heterocyclic ring; or

$\text{R}_a$ ,  $\text{R}_b$  and  $\text{R}_c$  together with the carbon atom to which they are attached  
form a tricyclic ring (for example adamantyl); or

$\text{R}_a$  and  $\text{R}_b$  are each independently  $(\text{C}_1\text{-C}_6)$ alkyl,  $(\text{C}_2\text{-C}_6)$ alkenyl,  $(\text{C}_2\text{-C}_6)$ alkynyl, phenyl $(\text{C}_1\text{-C}_6)$ alkyl, or a group as defined for  $\text{R}_c$  below other

hydrogen, -OH, -SH, halogen, -CN, -CO<sub>2</sub>H, (C<sub>1</sub>-C<sub>4</sub>)perfluoroalkyl, -CH<sub>2</sub>OH, -CO<sub>2</sub>(C<sub>1</sub>-C<sub>8</sub>)alkyl, -O(C<sub>1</sub>-C<sub>8</sub>)alkyl, -O(C<sub>2</sub>-C<sub>8</sub>)alkenyl, -S(C<sub>1</sub>-C<sub>8</sub>)alkyl, -SO(C<sub>1</sub>-C<sub>8</sub>)alkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>8</sub>)alkyl, -S(C<sub>2</sub>-C<sub>8</sub>)alkenyl, -SO(C<sub>2</sub>-C<sub>8</sub>)alkenyl, -SO<sub>2</sub>(C<sub>2</sub>-C<sub>8</sub>)alkenyl or a group -Q-W wherein Q represents a bond or -O-, -S-, -SO- or -SO<sub>2</sub>- and W represents a phenyl, phenylalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylalkyl, (C<sub>4</sub>-C<sub>8</sub>)cycloalkenyl, (C<sub>4</sub>-C<sub>8</sub>)cycloalkenylalkyl, heteroaryl or heteroarylalkyl group, which group W may optionally be substituted by one or more substituents independently selected from, hydroxyl, halogen, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub>(C<sub>1</sub>-C<sub>8</sub>)alkyl, -CONH<sub>2</sub>, -CONH(C<sub>1</sub>-C<sub>8</sub>)alkyl, -CONH(C<sub>1</sub>-C<sub>8</sub>)alkyl)<sub>2</sub>, -CHO, -CH<sub>2</sub>OH, (C<sub>1</sub>-C<sub>4</sub>)perfluoroalkyl, -O(C<sub>1</sub>-C<sub>8</sub>)alkyl, -S(C<sub>1</sub>-C<sub>8</sub>)alkyl, -SO(C<sub>1</sub>-C<sub>8</sub>)alkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>8</sub>)alkyl, -NO<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1</sub>-C<sub>8</sub>)alkyl, -N((C<sub>1</sub>-C<sub>8</sub>)alkyl)<sub>2</sub>, -NHCO(C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>4</sub>-C<sub>8</sub>)cycloalkenyl, phenyl or benzyl.

Examples of particular R<sub>3</sub> groups include methyl, ethyl, benzyl, 4-chlorobenzyl, 4-hydroxybenzyl, phenyl, cyclohexyl, cyclohexylmethyl, pyridin-3-ylmethyl, tert-butoxymethyl, naphthylmethyl, iso-butyl, sec-butyl, tert-butyl, 1-benzylthio-1-methylethyl, 1-methylthio-1-methylethyl, 1-mercapto-1-methylethyl, 1-methoxy-1-methylethyl, 1-hydroxy-1-methylethyl, 1-fluoro-1-methylethyl, hydroxymethyl, 2-hydroxyethyl, 2-carboxyethyl, 2-methylcarbamoyl, 2-carbamoyl, and 4-aminobutyl. Presently preferred R<sub>3</sub> groups include tert-butyl.

R<sub>4</sub> may be, for example

optionally substituted C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl or cycloalkyl;

optionally substituted phenyl, biphenyl or naphthyl;

optionally substituted piperidinyl, cyclohexyl, cyclopentyl, oxazolyl, thiazepinyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl,

optionally substituted piperidinyl, cyclohexyl, cyclopentyl, oxazolyl, thiazepinyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl, thienyl, furanyl, pyrrolyl, imidazolyl, benzofuranyl, benzothienyl, benzimidazolyl, thiazolyl, benzothiazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrazinyl, or triazinyl.

phenyl(C<sub>1</sub>-C<sub>6</sub> alkyl)-, phenyl(C<sub>2</sub>-C<sub>6</sub> alkenyl)- or phenyl(C<sub>2</sub>-C<sub>6</sub> alkynyl)- optionally substituted in the phenyl ring;

cycloalkyl(C<sub>1</sub>-C<sub>6</sub> alkyl)-, cycloalkyl(C<sub>2</sub>-C<sub>6</sub> alkenyl)- or cycloalkyl(C<sub>2</sub>-C<sub>6</sub> alkynyl)- optionally substituted in the cycloalkyl ring;


heterocyclyl(C<sub>1</sub>-C<sub>6</sub> alkyl)-, heterocyclyl(C<sub>2</sub>-C<sub>6</sub> alkenyl)- or heterocyclyl(C<sub>2</sub>-C<sub>6</sub> alkynyl)- optionally substituted in the heterocyclyl ring.

Specific examples of R<sub>4</sub> groups include

Methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 2-phenylcycloprop-1-yl, benzyl, biphenyl-2-yl, biphenyl-3-yl, biphenyl-4-yl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2,3-dimethoxyphenyl, 2,4-dimethoxyphenyl, 3,4-dimethoxyphenyl, 2-phenoxyphenyl, 3-phenoxyphenyl, 4-phenoxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2-iodophenyl, 3-iodophenyl, 4-iodophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 3,4-dimethyl, 2-aminophenyl, 3-aminophenyl, 4-aminophenyl, 2-acetamidophenyl, 3-acetamidophenyl, 4-acetamidophenyl, 2-N,N-dimethylaminophenyl, 3-N,N-dimethylaminophenyl, 4-N,N-dimethylaminophenyl, 2-methanesulfonamidophenyl, 3-methanesulfonamidophenyl, 4-methanesulfonamidophenyl, 2-hydroxyphenyl, 3-hydroxyphenyl, 4-hydroxyphenyl, 2,3-dihydroxyphenyl, 2,4-

dihydroxyphenyl, 3,4-dihydroxyphenyl, 2-thiophenolyl, 3-thiophenolyl, 4-thiophenolyl, 2-thioanisolyl, 3-thioanisolyl, 4-thioanisolyl, 1-naphthyl, 2-naphthyl, furan-2-yl, thien-2-yl, pyrrol-2-yl, 1-methylpyrrol-2-yl, imidazol-2-yl, 1-methylimidazol-2-yl, thiazol-2-yl, 5-phenylpyrrol-2-yl, 5-phenylfuran-2-yl, 5-phenylthien-2-yl, benzothiazol-2-yl, 1,2,4-oxadiazol-5-yl, 3-methyl-1,2,4-oxadiazol-5-yl, 3-phenyl-1,2,4-oxadiazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,2,4-thiadiazol-5-yl, 1,3,4-thiadiazol-2-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, N-oxides of pyridin-2-yl pyridin-3-yl and pyridin-4-yl, indol-2-yl, indol-3-yl, 1-methylindol-2-yl, 1-methylindol-3-yl, benzimidazol-2-yl, 1-methylbenzimidazol-2-yl, pyrazin-2-yl, 1,2-pyridazin-3-yl, 1,3-pyrimidin-2-yl, benzo[b]thien-2-yl, benzo[b]thien-3-yl, benzo[b]furan-2-yl, benzo[b]furan-3-yl, isoxazol-5-yl, quinolin-2-yl, quinolin-3-yl, isoquinolin-2-yl, isoquinolin-3-yl, 2-oxo-2-phenylethyl, diphenylmethyl, 4-N-methylaminophenyl, 4-N,N-dimethylcarboxamidophenyl, and 4-carboxyphenyl. R<sub>4</sub> may be a phenyl group which is substituted, for example in the 4-position, by one of the following:






In the compounds of formula (I) as defined above wherein Z is a radical of formula -C(=O)NH(OH) the radicals  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  may be any of those discussed above in relation to compounds (I) wherein Z is a radical of formula -N(OH)CH(=O). However, in addition,  $R_1$  may be, for example, a hydroxy, methoxy, ethoxy, n-propyloxy, allyloxy, amino, methylamino, dimethylamino, ethylamino, or diethylamino group.

Specific examples of compounds of the invention include those disclosed in the Examples herein.

Compounds of the invention wherein Z is a radical of formula -N(OH)CH(=O) may be prepared by N-formylation of a compound of formula (II)



wherein -OP represents a protected hydroxy group, and R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> are as defined in relation to formula (I) except that any reactive groups present therein which are reactive with the formylation reagent are protected, and then converting -OP to a hydroxy group and removing any protecting groups present in R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> or R<sub>4</sub>. Formylacetic anhydride is a convenient formylating agent for this reaction.

Compounds of formula (II) may be synthesised from easily accessible or commercially available starting materials, or by standard synthetic routes analogous to those shown in Schemes 1 and 2 below.

Hydroxamate compounds of formula (I) for use in accordance with the invention may be prepared by reacting the parent compound wherein Z is a carboxylic acid group with hydroxylamine or an N- and/or O-protected hydroxylamine, and thereafter removing any O- or N-protecting groups.

Antibacterial or antiprotozoal compositions with which the invention is concerned may be prepared for administration by any route consistent with the pharmacokinetic properties of the active ingredient(s).

Orally administrable compositions may be in the form of tablets, capsules, powders, granules, lozenges, liquid or gel preparations, such as oral, topical, or sterile parenteral solutions or suspensions. Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinyl-pyrrolidone; fillers for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricant, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants for example potato starch, or

acceptable wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, glucose syrup, gelatin hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

For topical application to the skin, the active ingredient(s) may be made up into a cream, lotion or ointment. Cream or ointment formulations which may be used for the drug are conventional formulations well known in the art, for example as described in standard textbooks of pharmaceuticals such as the British Pharmacopoeia.

The active ingredient(s) may also be administered parenterally in a sterile medium. Depending on the vehicle and concentration used, the drug can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle. Intra-venous infusion is another route of administration for the compounds used in accordance with the invention.

Safe and effective dosages for different classes of patient and for different disease states will be determined by clinical trial as is required in the art. It will be understood that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate

of excretion, drug combination and the severity of the particular disease undergoing therapy.

The following example illustrate embodiments of the invention.

The following abbreviations have been used throughout:


|      |  |
|------|--|
| CDI  | 1,1'-Carbonyldiimidazole                                     |
| DMF  | N,N-Dimethylformamide  |
| EDC  | N-Ethyl-N'-(3-dimethylaminopropyl)carbodiimide hydrochloride |
| TFA  | Trifluoroacetic acid   |
| HOBt | 1-Hydroxybenzotriazole                                       |
| HPLC | High performance liquid chromatography                       |
| LRMS | Low resolution mass spectrometry                             |
| THF  | Tetrahydrofuran  |
| TLC  | Thin layer chromatography                                    |

$^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded using a Bruker AC 250E spectrometer at 250.1 and 62.9 MHz, respectively. Mass spectra were obtained using a Perkin Elmer Sciex API 165 spectrometer using both positive and negative ion modes.

Where MIC values are quoted for the compounds of the examples, those results were obtained by the method described in the Biological Example.

### Example 1


2(R)-[[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(1'(S)-(4-fluorobenzoyl)-



2',2'-dimethylpropyl)amide.

The title compound was prepared as described in **Scheme 1**. The experimental details of the synthesis and relevant spectroscopic data are described below.

**Scheme 1**



**Reagents and Conditions:**

- A. Di-*tert*-butyldicarbamate, Et<sub>3</sub>N, MeOH
- B. HN(OMe)Me.HCl, CDI, CH<sub>2</sub>Cl<sub>2</sub>
- C. 4-Fluorophenylmagnesium bromide, THF, 0 °C to 25 °C, 20 hours
- D. TFA, CH<sub>2</sub>Cl<sub>2</sub>, <5 °C, 16 hours
- E. 2(R)-(Benzyloxy-amino-methyl)-hexanoic acid, WSC, 10 mol% HOBT, DMF, 16 hours
- F. Formylacetic anhydride, CH<sub>2</sub>Cl<sub>2</sub>, RT, 1 hour
- G. H<sub>2</sub>, 10% Pd/C, EtOAc/MeOH

**STEP A: 2(S)-tert-Butoxycarbonylamino-3,3-dimethyl-butyric acid**

To a suspension of L-*tert*-leucine (26.1 g, 0.2 mol) in methanol (150 mL) was added triethylamine (56 mL, 0.4 mol) and the mixture cooled to 0 °C. To the mixture was added slowly a solution of di-*tert*-butyldicarbamate (48 g, 0.22 mol) in methanol (40 mL) such that an internal temperature of between 0 and 5 °C was maintained. The reaction was allowed to stir overnight and the solvents were removed *in vacuo*. The residue was dissolved in ethyl acetate (200 mL) and washed with 10% w/v aqueous citric acid solution (3 x 100 mL). The organic layer was dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give the title product as a pale yellow oil (48.9 g, >100 %, residual solvent).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 9.20 (1H, bs), 5.10 (1H, m), 4.15 (1H, m), 1.45 (9H, s), 1.00 (9H, s).

**STEP B: [1(S)-(Methoxy-methyl-carbamoyl)-2,2-dimethylpropyl]-carbamic acid-*tert*-butyl ester.**

To a solution of 2(S)-*tert*-butoxycarbonylamino-3,3-dimethyl-butyric acid (30 g, 0.13 mol) in dichloromethane (400 mL) was added portionwise 1,1'-carbonyldiimidazole (63.3 g, 0.39 mol). After addition was complete the reaction was allowed to stir for 15 minutes before *N*-methoxy-*N*-methylamine hydrochloride (38 g, 0.39 mol) was added and the reaction allowed to stir for 16 hours. The solvents were removed *in vacuo* and the residue partitioned between ethyl acetate (250 mL) and 1M HCl (150 mL). The organic phase was washed with 1M HCl solution (2 x 100 mL), saturated sodium bicarbonate solution (2 x 100 mL) and brine (100 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 30% ethyl acetate in hexanes) afforded the title product as a pale yellow oil (26.5 g, 74 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 5.20 (1H, bd, J = 9.5 Hz), 4.65 (1H, d, J = 10.4 Hz), 3.77 (3H, s), 3.21 (3H, s), 1.43 (9H, s), 0.97 (9H, s).

LRMS: (m/z) 297 (M+Na)<sup>+</sup>.

**STEP C:** (1(S)-(4-Fluorobenzoyl)l-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester

[1(S)-(Methoxy-methyl-carbamoyl)-2,2-dimethylpropyl]-carbamic acid-*tert*-butyl ester (548 mg, 2.0 mmol) was dissolved in freshly distilled THF (10 mL) under an argon atmosphere. The solution was cooled to 0 °C whereupon 4-fluorophenylmagnesium bromide (8.0 mL, 1.0 M in THF, 8.0 mmol, 4 eqv) was added dropwise maintaining an internal temperature < 5 °C. The reaction was stirred at 0 °C for 30 minutes before warming to room temperature for a further 20 hours. After this time the reaction was cooled to 0 °C and saturated aqueous ammonium chloride solution (30 mL) was added. The aqueous phase was extracted with ethyl acetate (3 x 30 mL) and the organic phases washed with water (3 x 50 mL) and brine (50 mL), dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 20 % ethyl acetate in hexanes) afforded the title compound as a white solid (447 mg, 72%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.03 (2H, m), 7.14(2H, t, J = 8.5 Hz), 5.41 (1H, bd, J = 9.6 Hz), 5.12 (1H, d, J = 9.6 Hz), 1.43 (9H, s), 0.92 (9H, s).

LRMS: (m/z) 332 (M+Na)<sup>+</sup>.

**STEP D:** 2(S)-Amino-1-(4-fluorophenyl)-3,3-dimethyl-butan-1-one.

To a solution of (1(S)-(4-fluorobenzoyl)l-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester (319 mg, 1.03 mmol) in dichloromethane (5 mL) at 0 °C was added trifluoroacetic acid (2 mL). The reaction was placed at <5 °C for 16 hours before the solvents were removed *in vacuo* to give a bright yellow oil. The residue was dissolved in methanol (30 mL) and was treated portionwise with Dowex 1X8-400 basic resin until pH 9. Filtration and removal of the solvents *in vacuo* gave the title

compound as a pale yellow oil (~1.0 g, > 100 %, residual water). Purification by column chromatography (SiO<sub>2</sub>, 5 % methanol in dichloromethane) gave the title compound as a colourless oil (190 mg, 88 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.10 (2H, m), 7.15 (2H, t, J = 8.4 Hz), 4.20 (1H, s), 1.70 (2H, bs), 0.93 (9H, s).

LRMS: (m/z) 210 (M+H)<sup>+</sup>.

**STEP E:** 2(R)-(N-Benzyloxyaminomethyl)-hexanoic acid-(1'(S)-(4-fluorobenzoyl)-2',2'-dimethylpropyl)amide

To a suspension of 2(R)-(N-benzyloxyamino-methyl)-hexanoic acid (237 mg, 0.91 mmol, 1.1 eqv) in DMF (5 mL) at 0 °C was added EDC (200 mg, 1.1 eqv) and 1-hydroxybenzotriazole (12 mg, 10 mol%). The mixture was allowed to stir at 0 °C for 45 minutes before a solution of 2(S)-amino-1-(4-fluorophenyl)-3,3-dimethyl-butan-1-one (179 mg, 0.83 mmol, 1 eqv) in DMF (2 mL) was added. The reaction was allowed to warm to room temperature and stirred for 16 hours. The solvents were removed *in vacuo* and the residue dissolved in ethyl acetate (30 mL) and 10% w/v aqueous citric acid (40 mL). The aqueous phase was extracted with ethyl acetate (3 x 20 mL) and the combined organic layers washed with saturated sodium hydrogen carbonate (70 mL) and brine (2 x 70 mL). The organic phases were dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 30 % ethyl acetate in hexanes) gave the title compound as a clear oil (160 mg, 42 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.07 (2H, m), 7.35 (5H, m), 7.13 (2H, t, J = 8.7 Hz), 6.99 (1H, d, J = 9.0 Hz), 5.52 (1H, d, J = 9.0 Hz), 4.75 (2H, s), 3.10 (2H, m), 2.52 (1H, m), 1.70 - 1.10 (6H, m), 0.94 (9H, s), 0.81 (3H, t, J = 6.6 Hz).

LRMS: (m/z) 443 (M+H)<sup>+</sup>

**STEP F:** 2(R)-[N-Benzyloxy-N-formylamino]-methyl]-hexanoic acid-(1'(S)-(4-fluorobenzoyl)-2',2'-dimethylpropyl)amide.



To a solution of 2(*R*)-(*N*-benzyloxyaminomethyl)-hexanoic acid-(1'(*S*)-(4-fluorobenzoyl)-2',2'-dimethylpropyl)amide (150 mg, 0.35 mmol) in dichloromethane (10 mL) at 0 °C was added formylacetic anhydride (80 µL, 0.88 mmol, 2.5 eqv) and the reaction allowed to stir at room temperature. After 2 hours the solvents were removed *in vacuo* and the residue azeotroped with toluene (3 x 30 mL). The product was placed under high vacuum to give the title compound as a pale yellow solid (145 mg, 88 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.20 – 7.85 (3H, bm), 7.40 (5H, m), 7.15 (2H, m), 6.31 (1H, d, J = 9.3 Hz), 5.42 (1H, d, J = 9.2 Hz), 5.10 – 4.70 (2H, bm), 3.90 – 3.60 (2H, bm), 2.58 (1H, m), 1.80 – 1.05 (6H, m), 0.89 (9H, s), 0.78 (3H, t, J = 6.8 Hz).

LRMS: (*m/z*) 471 (M+H)<sup>+</sup>.

**STEP G:** 2(*R*)-[*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-fluorobenzoyl)-2',2'-dimethylpropyl)amide

To a solution 2(*R*)-[*N*-benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-(4-fluorobenzoyl)-2',2'-dimethylpropyl)amide (145 mg, 0.31 mmol) in methanol (10 mL) under an argon atmosphere was added a slurry of 10% Pd/C (15 mg) in ethyl acetate (1 mL). Hydrogen gas was bubbled through the reaction for 15 minutes after which time the reaction was allowed to stir under 1 atmosphere of hydrogen for two hours. The reaction was flushed with argon and filtered through glass wool. The solvents were removed *in vacuo* to afford the title compound as a hygroscopic white crystalline solid (105 mg, 90%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.40 (0.3H, s), 8.05 (2H, dd, J = 6.8, 5.4Hz), 7.86 (0.7H, s), 7.16 (2H, t, J = 3.2Hz), 6.90 (0.3H, bd, J = 9.1Hz), 6.82 (0.7H, bd, J = 9.2Hz), 5.50 (1H, d, J = 9.3Hz), 4.02 (0.2H, dd, J = 14.5, 7.5Hz), 3.87 (0.8H, dd, J = 14.2, 10.0Hz), 3.62 (0.2H, dd, J = 11.0, 3.0Hz), 3.46 (0.8H, dd, J = 14.1, 4.6Hz), 2.88-

2.70 (1H, m), 1.55 (1H, m), 1.45-1.10 (5H, m), 0.94 (2H, s), 0.90 (7H, s), 0.78 (3H, t, J = 6.6Hz).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ/ppm 199.6, 199.3, 173.5, 168.5, 164.5, 134.5, 131.8, 131.7, 116.7, 116.6, 116.3, 116.2, 59.5, 52.6, 48.4, 48.1, 45.0, 35.7, 30.5, 29.8, 29.5, 28.7, 28.5, 27.4, 22.9, 14.2.


LRMS: (m/z) 381 (M+H)<sup>+</sup>, 403 (M+Na)<sup>+</sup>.

MIC (*E. Coli*): 12.5 μM

MIC (*S. Capitis*): 25 μM

### Example 2

2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(1'(S)-(4-methoxybenzoyl 2',2'dimethylpropyl)amide



The title compound was prepared as described in **Scheme 1** in an analogous manner to 2(R)-[N-formyl-N-hydroxyamino)-methyl]-hexanoic acid-(1'(S)-(4-fluorobenzoyl)-2',2'-dimethylpropyl)amide . The relevant spectroscopic data are described below.

**STEP C:** (1(S)-(4-Methoxybenzoyl)-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester  
Purification by column chromatography (SiO<sub>2</sub>, 20 % ethyl acetate in hexanes)  
afforded the title compound as a white solid (836 mg, 71%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.00 (2H, d, J = 8.8 Hz), 6.95 (2H, d, J = 8.8 Hz), 5.45 (1H, d, J = 9.5 Hz), 5.12 (1H, d, J = 9.8 Hz), 3.89 (3H, s), 1.43 (9H, s), 0.93 (9H, s).

**STEP D:** 2(*S*)-Amino-1-(4-methoxyphenyl)-3,3-dimethyl-butan-1-one.

Purification by column chromatography (SiO<sub>2</sub>, 5 % methanol in dichloromethane) gave the title compound as a colourless oil (176 mg, 78 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.95 (2H, d, J = 9.0 Hz), 6.93 (2H, d, J = 9.0 Hz), 4.20 (1H, s), 3.89 (3H, s), 1.68 (2H, bs), 0.94 (9H, s).

LRMS: (*m/z*) 222 (M+H)<sup>+</sup>.

**STEP E:** 2(*R*)-(N-Benzyloxyaminomethyl)-hexanoic acid-(1'(*S*)-(4-methoxybenzoyl)-2',2'-dimethylpropyl)amide

Purification by column chromatography (SiO<sub>2</sub>, 30 % ethyl acetate in hexanes) gave the title compound as a clear oil (215 mg, 61 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.01 (2H, d, J = 9.0 Hz), 7.35 (5H, m), 6.95 (2H, d, J = 9.0 Hz), 5.55 (1H, d, J = 9.5 Hz), 4.73 (2H, m), 3.87 (3H, s), 3.10 (2H, m), 2.51 (1H, m), 1.70 – 1.10 (6H, m), 0.94 (9H, s), 0.80 (3H, t, J = 6.7 Hz).

LRMS: (*m/z*) 455 (M+H)<sup>+</sup>

**STEP F:** 2(*R*)-[N-Benzyloxy-N-formylamino)-methyl]-hexanoic acid-(1'(*S*)-(4-methoxybenzoyl)-2',2'-dimethylpropyl)amide

The title compound was isolated as a pale yellow solid (187 mg, 88 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.20 – 7.85 (3H, m), 7.60 – 7.30 (5H, m), 7.05 – 6.90 (2H, bd, J = 8.7 Hz), 6.40 (1H, bd, J = 9.2 Hz), 5.50 (1H, bd, J = 9.3 Hz), 5.10 – 4.70 (2H,

bm), 3.92 (3H, bs), 3.75 (2H, bm), 2.60 (1H, m), 1.90 – 1.10 (6H, m), 0.89 (9H, s), 0.77 (3H, m).

LRMS: ( $m/z$ ) 483 (M+H)<sup>+</sup>.

**STEP G:** 2(*R*)-[*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-methoxybenzoyl)-2',2'-dimethylpropyl)amide

The title compound was isolated as a hygroscopic white crystalline solid (133 mg, 98%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.40 (0.3H, s), 8.00 (2H, d, J = 8.8Hz), 7.86 (0.7H, s), 6.95 (2H, d, J = 8.8Hz), 6.76 (1H, d, J = 9.3Hz), 5.52 (1H, d, J = 9.4Hz), 4.10 (1H, m), 3.88 (3H, s), 3.62 (0.3H, dd, J = 14.6, 3.1Hz), 3.47 (0.7H, dd, J = 14.0, 4.0Hz), 2.85 (0.7H, m), 2.72 (0.3H, m), 1.55 (1H, m), 1.45-1.10 (5H, m), 0.94 (3H, s), 0.90 (6H, s), 0.77 (3H, t, J = 6.6Hz).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ/ppm 199.2, 198.9, 176.3, 173.3, 164.7, 164.5, 131.6, 131.5, 130.7, 114.7, 114.4, 59.3, 59.0, 55.9, 52.5, 48.3, 46.5, 45.1, 36.1, 35.9, 30.4, 30.3, 29.6, 27.5, 22.9, 14.2.


LRMS: ( $m/z$ ) 391 (M-H)<sup>-</sup>; 393 (M+H)<sup>+</sup>, 415 (M+Na)<sup>+</sup>.

MIC (*E. Coli*): 12.5 μM

MIC (*S. Capitis*): 6.25 μM

**Example 3**

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-cyclohexanecarbonyl-2'-methylpropyl)amide



The title compound was prepared in an analogous manner to that described in **Scheme 1** utilising 2(*S*)-*tert*-butoxycarbonylamino-3-methyl-butyric acid as the starting material in **Step B**. The relevant spectroscopic data are described below.

**STEP B:** [1(*S*)-(Methoxy-methyl-carbamoyl)-2-methylpropyl]-carbamic acid-*tert*-butyl ester.

The title product was isolated as a pale yellow oil (14.7 g, 82 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 5.13 (1H, m), 4.57 (1H, m), 3.77 (3H, s), 3.21 (3H, s), 1.99 (1H, m), 1.44 (9H, s), 0.96 (3H, d, J = 6.8 Hz), 0.90 (3H, d, J = 6.8 Hz).

**STEP C:** (1(*S*)-(Cyclohexanecarbonyl)-2-methylpropyl)carbamic acid-*tert*-butyl ester

Purification by column chromatography (SiO<sub>2</sub>, 25 % ethyl acetate in hexanes) afforded the title compound as a white solid (253 mg, 42%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 5.13 (1H, bd, J = 8.5 Hz), 4.40 (1H, m), 2.52 (1H, m), 2.13 (1H, m), 2.00 – 1.60 (7H, m), 1.44 (9H, s), 1.25 (3H, m), 1.01 (3H, d, J = 6.9 Hz), 0.75 (3H, d, J = 6.9 Hz).

**STEP D:** 2(*S*)-Amino-1-cyclohexyl-3-methyl-butan-1-one

The title compound was isolated as colourless oil (158 mg, 98 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 3.49 (1H, d, J = 3.7 Hz), 2.55 (1H, m), 2.12 (1H, m), 1.95 - 1.10 (10H, m), 1.03 (3H, d, J = 6.9 Hz), 0.75 (3H, d, J = 6.8 Hz).

**STEP E:** 2(*R*)-(*N*-Benzyloxyaminomethyl)-hexanoic acid-(1'(*S*)-cyclohexanecarbonyl-2'-methylpropyl)amide

Purification by column chromatography (SiO<sub>2</sub>, 25 % ethyl acetate in hexanes) gave the title compound as a clear oil (116 mg, 35 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.35 (5H, m), 6.49 (1H, d, J = 8.9 Hz), 5.70 (1H, bs), 4.78 (1H, dd, J = 8.9, 4.7 Hz), 4.72 (2H, s), 3.08 (2H, m), 2.52 (2H, m), 2.15 (1H, m), 2.05 - 1.10 (16H, m), 0.98 (3H, d, J = 6.8 Hz), 0.86 (3H, t, J = 6.9 Hz), 0.78 (3H, d, J = 6.8 Hz).

LRMS: (*m/z*) 417 (M+H)<sup>+</sup>

**STEP F:** 2(*R*)-[*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-cyclohexanecarbonyl-2'-methylpropyl)amide

The title compound was isolated as a viscous oil (105 mg, 99 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.30 - 7.80 (1H, m), 7.40 (5H, m), 6.12 (1H, m), 5.15 - 4.60 (3H, m), 3.95 - 3.50 (2H, m), 3.10 (1H, m), 2.55 (2H, m), 1.10 - 2.30 (16H, m), 1.10 - 0.60 (9H, m).

LRMS: (*m/z*) 445 (M+H)<sup>+</sup>, 467 (M+Na)<sup>+</sup>.

**STEP G:** 2(*R*)-[*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-cyclohexanecarbonyl-2'-methylpropyl)amide

The title compound was isolated as a hygroscopic white crystalline solid (67 mg, 84 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.37 (0.3H, s), 7.81 (0.7H, s), 6.69 (1H, d, J = 9.0Hz), 4.74 (1H, dd, J = 9.0, 5.0Hz), 4.00 (0.3H, dd, J = 14.6, 7.6Hz), 3.84 (0.7H, dd, J = 14.0, 9.8Hz), 3.59 (0.3H, dd, J = 14.5, 3.4Hz), 3.43 (0.7H, dd, J = 14.1, 3.7Hz), 2.83 (0.7H, m), 2.70 (0.3H, m), 2.50 (1H, m), 2.12 (1H, m), 2.00-1.10 (16H, m), 1.00-0.60 (9H, m).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 176.3, 173.5, 61.5, 61.3, 52.9, 49.5, 49.3, 48.5, 30.7, 30.4, 30.2, 30.0, 29.8, 29.6, 27.6, 26.3, 26.1, 25.6, 25.5, 23.0, 20.3, 17.6, 14.3.


LRMS: (m/z) 355 (M+H)<sup>+</sup>, 377 (M+Na)<sup>+</sup>.

MIC (*E. Coli*): 50 μM

MIC (*S. Capitis*): 25 μM

#### Example 4

3-Cyclopentyl-2(*R*)-[(*N*-formyl-*N*-hydroxyamino)-methyl]-*N*-[1'(*S*)-(4-fluorobenzoyl)-2',2'-dimethylpropyl]propionamide



The title compound was prepared in an analogous manner to that described in **Scheme 1** utilising 2(*R*)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid in **Step E**. Spectroscopic data are described below. Experimental details of the synthesis of 2(*R*)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid and relevant spectroscopic data are also presented.

**STEP G: 3-Cyclopentyl-2(R)-[(N-formyl-N-hydroxyamino)-methyl]-N-[1'(S)-(4-fluorobenzoyl)-2',2'-dimethylpropyl]propionamide**

The title compound was isolated as a hygroscopic white crystalline solid (120 mg, 81 %).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):** δ/ppm 8.27 (0.3H, s), 8.10 (2H, m), 7.85 (0.7H, s), 7.23 (2H, t, J=8.7Hz), 5.47 (1H, m), 3.77 (1H, m), 3.58 (0.3H, dd, J=14.0, 5.6Hz), 3.42 (0.7H, dd, J=14.1, 4.7Hz), 3.09 (0.7H, m), 2.93 (0.3H, m), 1.82-1.20 (9H, m), 1.15-0.80 (11H, m).

**<sup>13</sup>C NMR (CD<sub>3</sub>OD):** δ/ppm 176.8, 176.5, 169.8, 165.7, 136.3, 132.9, 132.8, 117.4, 117.0, 61.1, 54.3, 45.1, 45.0, 39.3, 38.1, 38.0, 36.2, 34.4, 33.7, 33.6, 27.8, 26.4, 26.3.


**LRMS:** (*m/z*) 407 (M+H)<sup>+</sup>.

**MIC (*E. Coli*):** 3.1 μM

**MIC (*S. Capitis*):** 3.1 μM

**Example 5**

**3-Cyclopentyl-2(R)-[(N-formyl-N-hydroxyamino)-methyl]-N-[1'(S)-(4-methoxybenzoyl)-2',2'-dimethylpropyl]propionamide**





The title compound was prepared in an analogous manner to that described in **Scheme 1** utilising 2(*R*)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid in **Step E**. Spectroscopic data are described below.

**STEP G:** 3-Cyclopentyl-2(*R*)-[(*N*-formyl-*N*-hydroxyamino)-methyl]-*N*-[1'(*S*)-(4-methoxybenzoyl)-2',2'-dimethylpropyl]propionamide

The title compound was isolated as a hygroscopic white crystalline solid (173 mg, 96 %).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):** δ/ppm 8.26 (0.4H, s), 8.01 (2H, d, J=9.0Hz), 7.85 (0.6H, s), 7.01 (2H, d, J=9.0Hz), 5.49 (1H, bs), 3.87 (3H, s), 3.79 (1H, m), 3.59 (0.4H, dd, J=11.7, 5.6Hz), 3.41 (0.6H, dd, J=14.4, 4.9Hz), 3.09 (0.6H, m), 2.92 (0.4H, m), 1.85-1.20 (9H, m), 1.15-0.80 (11H, m).


**<sup>13</sup>C NMR (CD<sub>3</sub>OD):** δ/ppm 200.0, 176.6, 176.4, 166.0, 132.5, 132.3, 115.4, 60.6, 60.5, 56.5, 54.3, 51.9, 45.3, 45.1, 39.3, 38.1, 36.3, 34.3, 33.8, 33.6, 27.9, 26.4, 26.3.

**LRMS:** (*m/z*) 419 (M+H)<sup>+</sup>.

**MIC (*E. coli*):** 3.1 μM

**MIC (*S. capitis*):** 1.6 μM

## Synthesis of 2(R)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid



Reagents and Conditions: 1:  $\text{TiCl}_4$ , trioxane,  $\text{CH}_2\text{Cl}_2$ ; 2:  $\text{H}_2\text{O}_2$ ,  $\text{LiOH}$ ; 3:  $\text{H}_2\text{NOBn}$ , WSC,  $\text{THF}/\text{H}_2\text{O}$ ; 4:  $\text{Ph}_3\text{P}$ , DIAD,  $\text{THF}$ ; 5:  $\text{LiOH}$ ,  $\text{THF}/\text{MeOH}/\text{H}_2\text{O}$ .

**Step 1: 4S-Benzyl-3-[3-cyclopentyl-2R-hydroxymethyl-propionyl]-oxazolidin-2-one**

To a stirred, cooled ( $0\text{ }^\circ\text{C}$ ) solution of 4S-benzyl-(3-cyclopentyl-propionyl)-oxazolidin-2-one (21 g, 69.8 mmol) in dichloromethane (350 ml) was added a solution of titanium tetrachloride (1M in dichloromethane, 73.25 ml, 73.2 mmol), dropwise. The resulting yellowish slurry was stirred for 10 minutes at  $0\text{ }^\circ\text{C}$ , and then DIPEA (13.37 ml, 76.7 ml) was added dropwise to furnish a dark-red solution. The stirring was maintained for 1 h at  $0\text{ }^\circ\text{C}$ , and then a solution of s-trioxane (7.53 g, 83.7 mmol), in dichloromethane (70 ml) was added dropwise followed by the addition of a solution of titanium tetrachloride (1M in dichloromethane, 73.25 ml, 73.2 mmol). The reaction mixture was then stirred for 4 h at  $0\text{ }^\circ\text{C}$ . Saturated aqueous ammonium chloride (250 ml) was added to the reaction mixture and the aqueous layer was extracted with additional dichloromethane (2x300 ml). The combined organic layers were washed with water (150 ml) and with brine (80 ml), dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to yield a yellow solid which on trituration with diethyl ether furnished a white solid (16.57 g, 72%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.38-7.22 (5H, m), 4.70 (1H, m), 4.22-4.18 (2H, m), 3.99 (1H, m), 3.96-3.75 (2H, m), 3.31 (1H, dd, J = 13.4 & 3.3 Hz), 2.82 (1H, dd, J = 13.4 & 9.4 Hz), 2.24 (1H, dd, J = 8.3 & 4.5 Hz), 2.81-1.30 (4H, m) and 1.13 (1H, m);  
<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ/ppm 176.3, 153.6, 135.2, 129.5, 129.0, 127.4, 66.2, 64.2, 55.7, 44.8, 37.9, 37.8, 34.6, 33.0, 32.4 and 25.1.

**Step 2: 3-Cyclopentyl-2R-hydroxymethyl-propionic acid**

To a stirred, cooled (0 °C) solution of 4S-Benzyl-3-[3-cyclopentyl-2R-hydroxymethyl-propionyl]-oxazolidin-2-one (16.05 g, 48.5 mmol) in THF-water (4:1, 250 ml) was added 27.5% aqueous hydrogen peroxide (24 ml, 194 mmol), followed by lithium hydroxide monohydrate (4.07 g, 97 mmol) in water (50 ml). After the reaction was complete (30 min), THF was removed *in vacuo*. The aqueous layer was extracted with dichloromethane (3x100 ml) and acidified to pH 2 with 4M hydrochloric acid. The aqueous layer was extracted with diethyl ether (2x150 ml). The combined organic layers were washed with brine (60 ml), dried over anhydrous magnesium sulfate and filtered. The solvent was removed *in vacuo* to afford a yellow oil which was further purified by column chromatography (25% ethyl acetate in hexanes to 100% ethyl acetate) to furnish the title compound as an oil (8.3 g, quant.).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 6.60-5.90 (1H, br s), 3.80-3.78 (2H, m), 2.67 (1H, m), 1.98-1.40 (9H, m) and 1.20-0.98 (2H, m).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ/ppm 181.0, 63.2, 46.9, 37.8, 34.5, 32.7, 32.6, 25.1 and 25.1.

**Step 3: N-Benzyl-3-cyclopentyl-2R-hydroxymethyl-propionamide**

To a stirred, cooled (0 °C) mixture of 3-cyclopentyl-2R-hydroxymethyl-propionic acid (1.1 g, 6.4 mmol) in THF-water (4:1, 30 ml), was added O-benzylhydroxylamine. The pH of the resulting solution was adjusted to 4.5 by addition of 1M hydrochloric acid, and then EDC (1.84 g, 9.6 mmol) was added in one portion. The resulting solution

was stirred for 2.5 h at room temperature while controlling pH at 4.5 by addition of 1M hydrochloric acid. After removal of the THF, the aqueous layer was extracted with ethyl acetate (3x40 ml) and the combined organic layers were washed with 10% citric acid (3x15 ml), 5% sodium hydrogen carbonate and dried over anhydrous magnesium sulfate. The solvent was removed *in vacuo* to afford the title compound as a colourless crystalline solid (1.18 g, 67%). This compound was then used without any further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.14 (1H, br s), 7.40-7.34 (5H, m), 4.94 (2H, br s), 3.76-3.66 (2H, m), 1.79-1.47 (11H, m) and 1.17-0.97 (2H, m).

LRMS: +ve ion 278 [M+H], 555 [2M+H].

#### Step 4: N-Benzyloxy-3R-cyclopentylmethyl-azetidin-2-one

To a stirred, cooled (0 °C) solution of *N*-Benzyloxy-3-cyclopentyl-2R-hydroxymethyl-propionamide (8.63 g, 31.1 mmol) and triphenylphosphine (9 g, 34.2 mmol) in dry THF (320 ml) was added DIAD (6.12 ml, 31.1 mmol), dropwise. The resulting solution was stirred at room temperature overnight. After removal of THF *in vacuo*, the residue was purified by column chromatography (hexanes:ethyl acetate, 5:1 to 3:1) to give the desired product as a white solid (6.7 g, 83%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.76-7.39 (5H, m), 4.94 (2H, br s), 3.36 (1H, m), 2.96-2.80 (2H, m), 1.89-1.38 (9H, m) and 1.18-0.98 (2H, m).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ/ppm 167.7, 129.6, 129.3, 129.0, 78.1, 52.5, 45.1, 39.1, 35.2, 33.1, 32.9, 25.5 and 25.3.

LRMS: +ve ion 260 [M+H], 519 [2M+H].

#### Step 5: 2R-(Benzyloxyamino-methyl)-3-cyclopentyl-propionic acid

To a stirred, cooled (0 °C) solution of *N*-Benzyloxy-3R-cyclopentylmethyl-azetidin-2-one (6.7 g, 25.8 mmol) in THF-methanol (3:1, 100 ml) was added lithium hydroxide

monohydrate (1.3 g, 31.0 mmol) in water (25 ml). The reaction mixture was stirred and allowed to warm to room temperature overnight. The solvent was removed *in vacuo* and the aqueous layer was extracted with diethyl ether, then acidified to pH 2 by addition of 4M hydrochloric acid. The aqueous layer was extracted with diethyl ether (3x40 ml), and the combined organic layers were washed with brine, dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to give the title compound as white crystals (6.02 g, 84%).


<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.68-7.30 (5H, m), 4.78-4.68 (2H, m), 3.12-3.10 (2H, d, J = 6.9 Hz), 2.76 (1H, m), 1.91-1.39 (11H, m), 1.20-1.00 (2H, m).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ/ppm 180.1, 137.7, 129.0, 128.9, 128.5, 78.0, 53.9, 42.9, 38.3, 36.6, 33.1, 33.0, 25.5.

LRMS: 276 [M-H], 553 [2M-H].

### Example 6

2(R)-[[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(S)-(1-methyl-1H-imidazole-2-carbonyl)-propyl]-amide.



The title compound was prepared as described in Scheme 2. The experimental details of the synthesis and relevant spectroscopic data are described below.

**STEP A:** 2(S)-Amino-N-methoxy-3,3,N-trimethyl-butylamide

To a solution of {1(*S*)-(methoxy-methyl-carbamoyl)-2,2-dimethylpropyl}-carbamic acid-*tert*-butyl ester (1.686 g, 6.15 mmol) in dichloromethane (20 mL) at 0 °C was added trifluoroacetic acid (17 mL). The reaction was allowed to stand for 16 hours at <5 °C before the solvents were removed *in vacuo* to give a bright yellow oil. The residue was dissolved in methanol (30 mL) and was treated portionwise with Dowex 1X8-400 basic resin until pH 9. Filtration and removal of the solvents *in vacuo* gave the title compound as a pale yellow oil (990 mg, 92 %). The material was used without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 3.70 (3H, s), 3.61 (1H, bs), 3.20 (3H, s), 1.60 (2H, bs), 0.98 (9H, s).

LRMS: (*m/z*) 175 (M+H)<sup>+</sup>.

**STEP B:** 2(*R*)-[(*N*-Benzyloxyamino)-methyl]-hexanoic acid-[1'(*S*)-(methoxy-methyl-carbamoyl)-2',2'-dimethylpropyl]amide

To a suspension of 2(*R*)-(*N*-benzyloxyamino-methyl)-hexanoic acid (1.430 g, 5.17 mmol, 1.1 eqv) in DMF (20 mL) at 0 °C was added EDC (1.190 g, 5.64 mmol, 1.2 eqv) and 1-hydroxybenzotriazole (84 mg, 10 mol%). The mixture was allowed to stir at 0 °C for 45 minutes before a solution of 2(*S*)-amino-*N*-methoxy-3,3,*N*-trimethylbutyramide (900 mg, 4.7 mmol, 1 eqv) in DMF (5 mL) was added. The reaction was allowed to warm to room temperature and stirred overnight. The solvents were removed *in vacuo* and the residue dissolved in ethyl acetate (75 mL) and 10% w/v aqueous citric acid (50 mL). The aqueous phase was extracted with ethyl acetate (3 x 75 mL) and the combined organic layers washed with saturated sodium hydrogen carbonate solution (100 mL) and brine (2 x 100 mL). The organic phases were dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 25% ethyl acetate in hexanes) gave the title compound as a clear oil (1.43 g, 68 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.35 (5H, m), 6.73 (1H, bd, J = 10.0 Hz), 5.70 (1H, bs), 5.05 (1H, d, J = 10.0 Hz), 4.74 (2H, s), 3.78 (3H, s), 3.20 (3H, s), 3.07 (2H, m), 2.50 (1H, m), 1.65 (1H, m), 1.45 (1H, m), 1.25 (4H, m), 0.98 (9H, s), 0.86 (3H, t, J = 7.0 Hz).  
LRMS: (m/z) 408 (M+H)<sup>+</sup>.

**STEP C:** 2(R)-[(N-Benzyloxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(S)-(1-methyl-1H-imidazole-2-carbonyl)-propyl]-amide.

To a solution of *N*-methyl-imidazole (246 mg, 3 mmol, 6 eqv), in dry freshly distilled THF (5 mL) under argon at -78 °C was added dropwise *n*-BuLi (1.84 mL, 1.6M in hexanes, 3.0 mmol, 6 eqv) such that an internal temperature of < -70 °C was maintained. The reaction was allowed to stir at -70 °C for 90 minutes after which time a solution of 2(R)-[(N-benzyloxyamino)-methyl]-hexanoic acid-[1'(S)-(methoxymethyl-carbamoyl)-2',2'-dimethylpropyl]amide (204 mg, 0.5 mmol, 1eqv) in freshly distilled THF (2 mL) was added dropwise such that an internal temperature of < -70 °C was maintained. The reaction was allowed to stir at -70 °C for five hours before saturated aqueous ammonium chloride solution (10 mL) was added and the solution allowed to warm to room temperature. The aqueous phase was extracted with ethyl acetate (3 x 50 mL), and the organic phases combined and washed with water (3 x 50 mL) and brine (50 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 30 % hexanes in ethyl acetate) gave the title product as a colourless oil (77 mg, 36 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.33 (5H, m), 7.17 (1H, s), 7.03 (1H, s), 5.75 (1H, bs), 5.65 (1H, d, J = 9.1 Hz), 4.75 (2H, s), 3.98 (3H, s), 3.10 (2H, m), 2.54 (1H, m), 1.90 (1H, bm), 1.58 (1H, m), 1.40 (1H, m), 1.23 (4H, m), 0.99 (9H, s), 0.80 (3H, t, J = 6.7 Hz).  
LRMS: (m/z) 429 (M+H)<sup>+</sup>.

**STEP D:** 2(R)-[(N-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(S)-(1-methyl-1H-imidazole-2-carbonyl)-propyl]-amide.

To a solution of 2(*R*)-[(*N*-benzyloxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(*S*)-(1-methyl-1*H*-imidazole-2-carbonyl)-propyl]-amide (77 mg, 0.18 mmol) in dichloromethane (2 mL) at 0 °C was added formylacetic anhydride (50  $\mu$ L, 0.54 mmol, 3 eqv) and the reaction allowed to stir at room temperature. After 2 hours the solvents were removed *in vacuo* and the residue azeotroped with toluene (2 x 30 mL). The product was placed under high vacuum for several hours to give the title compound as an off white waxy solid (76 mg, 93 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm 8.30 – 7.80 (1H, m), 7.40 (5H, bm), 7.19 (1H, bs), 7.05 (1H, bs), 5.55 (1H, bd, *J* = 9.3 Hz), 5.15 – 4.70 (2H, bm), 3.96 (3H, bs), 3.80 (1H, bm), 3.15 (1H, bm), 2.60 (1H, m), 1.70 – 0.70 (18H, m).

LRMS: (*m/z*) 456 (M+H)<sup>+</sup>.

**STEP E:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(*S*)-(1-methyl-1*H*-imidazole-2-carbonyl)-propyl]-amide.

To a solution of 2(*R*)-[(*N*-benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(*S*)-(1-methyl-1*H*-imidazole-2-carbonyl)-propyl]-amide. (76 mg, 0.17 mmol) in methanol (2 mL) was added a slurry of 10% Pd/C (10 mg) in ethyl acetate (1 mL). Hydrogen gas was bubbled through the reaction for 15 minutes after which time the reaction was allowed to stir under one atmosphere of hydrogen for 2 hours. The reaction was flushed with argon and filtered through glass wool. The solvents were removed *in vacuo* to afford the title compound as a hygroscopic white crystalline solid (43 mg, 70%).

<sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$ /ppm 8.26 (0.3H, s), 8.22 (0.7H, d, *J* = 9.0Hz), 8.10 (0.3H, d, *J* = 8.0Hz), 7.84 (0.7H, s), 7.35 (1H, s), 7.14 (1H, s), 5.71 (1H, m), 3.99 (3H, s), 3.72 (1H, m), 3.40 (0.5H, dd, *J* = 14.1, 4.6Hz), 3.31 (0.5H, m), 3.05 (0.7H, m), 2.88 (0.3H, m), 1.60-1.10 (6H, m), 0.99 (4H, s), 0.97 (5H, s), 0.82 (3H, t, *J* = 6.6Hz).



$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta/\text{ppm}$  194.7, 178.7, 147.2, 132.9, 132.4, 65.2, 56.5, 48.2, 47.9, 39.6, 39.1, 34.1, 33.2, 30.4, 26.6 and 17.2.


LRMS: ( $m/z$ ) 365 ( $\text{M}-\text{H}$ ) $^-$ , 367 ( $\text{M}+\text{H}$ ) $^+$ .

MIC (*E. coli*): 25  $\mu\text{M}$

MIC (*S. capitis*): 200  $\mu\text{M}$

### Example 7

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(*S*)-(pyridine-2-carbonyl)-propyl]-amide



The title compound was prepared from 2-bromopyridine as outlined in **Scheme 2**, in an analogous manner to 2(*R*)-[(*N*-formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(*S*)-(1-methyl-1*H*-imidazole-2-carbonyl)-propyl]-amide. The relevant spectroscopic data are described below.

**STEP C:** 2(*R*)-[(*N*-Benzyloxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(*S*)-(pyridine-2-carbonyl)-propyl]-amide.

Purification by column chromatography ( $\text{SiO}_2$ , 30 % to 50 % ethyl acetate in hexanes) gave the title product as a colourless oil (160 mg, 60 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.72 (1H, m), 8.05 (1H, m), 7.82 (1H, m), 7.45 (1H, m), 7.40 – 7.28 (5H, m), 7.00 (1H, bd, J = 9.3 Hz), 6.22 (1H, d, J = 9.3 Hz), 5.74 (1H, bs), 4.76 (2H, s), 3.11 (2H, m), 2.52 (1H, m), 1.58 (1H, m), 1.40 (1H, m), 1.22 (4H, m), 0.95 (9H, s), 0.81 (3H, t, J = 6.6 Hz).

LRMS: (m/z) 426 (M+H)<sup>+</sup>.

**STEP D:** 2(R)-[(N-Benzyloxy-N-formylamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(S)-(pyridine-2-carbonyl)-propyl]-amide.

The title compound was isolated as a pale yellow waxy solid (162 mg, 90 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.75 (1H, m), 8.35 – 7.80 (4H, m), 7.60 – 7.10 (5H, m), 6.10 (1H, m), 4.90 (2H, m), 3.75 (2H, bm), 2.90 (2H, bm), 1.60 – 1.10 (6H, bm), 0.90 (9H, s), 0.85 (3H, m).

LRMS: (m/z) 454 (M+H)<sup>+</sup>.

**STEP E:** 2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-[2,2-dimethyl-1(S)-(pyridine-2-carbonyl)-propyl]-amide.

The title compound was isolated as a hygroscopic white crystalline solid (113 mg, 94 %).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.72 (1H, m), 8.27 (0.3H, s), 8.10-7.90 (2H, m), 7.85 (0.7H, s), 7.59 (1H, m), 7.40-7.20 (1H), 6.18 (1H, m), 3.85-3.58 (1.7H, m), 3.41 (0.3H, dd, J = 14.2, 4.7Hz), 3.05 (0.7H, m), 2.89 (0.3H, m), 1.60-1.05 (6H, m), 0.97 (3.5H, s), 0.95 (5.5H, s), 0.79 (3H, t, J = 6.7Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 176.5, 176.2, 155.3, 150.6, 139.1, 129.2, 123.9, 122.3, 60.1, 53.5, 45.6, 45.3, 35.8, 32.2, 30.7, 27.9, 24.1, 14.8.


LRMS: (m/z) 364 (M+H)<sup>+</sup>.

MIC (*E. Coli*): 12.5 μM

MIC (*S. Capitis*): 50  $\mu$ M

### Example 8

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(benzofuran-2-carbonyl)-2',2'-dimethylpropyl)amide.



The title compound was prepared from 2-lithiobenzofuran as outlined in **Scheme 2**, in an analogous manner to 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-[2',2'-dimethyl-1'(*S*)-(1-methyl-1*H*-imidazole-2-carbonyl)-propyl]-amide. The relevant spectroscopic data are described below.

**STEP C:** 2(*R*)-[(*N*-Benzyloxyamino)-methyl]-hexanoic acid-[1'(*S*)-(benzofuran-2-carbonyl)-2',2'-dimethyl-propyl]-amide.

Purification by column chromatography (SiO<sub>2</sub>, 25% ethyl acetate in hexanes) gave the title compound as a colourless oil (43 mg, 19 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.90 – 7.20 (10H, m), 7.05 (1H, bd, *J* = 9.2 Hz), 5.72 (1H, bs), 5.45 (1H, d, *J* = 9.2 Hz), 4.76 (2H, s), 3.11 (2H, m), 2.53 (1H, m), 1.80 – 1.10 (6H, m), 1.03 (9H, s), 0.82 (3H, m).

LRMS: (*m/z*) 465 (M+H)<sup>+</sup>.

**STEP D:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-[1'(*S*)-(bezofuran-2-carbonyl)-2',2'-dimethyl-propyl]-amide.

Purification by column chromatography (SiO<sub>2</sub>, 2:1 ethyl acetate : hexanes) gave the title compound as a clear oil (24 mg, 52 %).

**<sup>1</sup>H NMR (MeOD):** 8.13 (0.5H, bs), 7.90 (0.5H, bs), 7.81 (2H, m), 7.70 – 7.25 (8H, m), 5.25 (1H, m), 4.90 (2H, m), 3.75 (2H, m), 2.90 (1H, m), 1.60 – 1.00 (6H, m), 0.98 (9H, s), 0.71 (3H, m).

**LRMS:** (*m/z*) 493 (M+H)<sup>+</sup>, 515 (M+Na)<sup>+</sup>.

**STEP E:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-[1(*S*)-(bezofuran-2-carbonyl)-2,2-dimethyl-propyl]-amide.

The title compound was isolated as a hygroscopic white crystalline solid (20 mg, 100%).

**<sup>1</sup>H NMR (MeOD):** 8.27 (0.3H, s), 7.85 (1.7H, m), 7.80 (1H, d, J=7.8Hz), 7.62 (1H, d, J=8.4Hz), 7.54 (1H, t, J=7.8Hz), 7.35 (1H, m), 5.35 (1H, br. s), 3.78 (1H, m), 3.62 (0.3H, dd, J=14.0, 5.6Hz), 3.42 (0.7H, dd, J=14.0, 4.7Hz), 3.09 (0.7H, m), 2.91 (0.3H, m), 1.42 (2H, m), 1.25 - 0.95 (13H, bm), 0.80-0.65 (3H, m).

**<sup>13</sup>C NMR (MeOD):** 191.5, 176.5, 157.7, 154.3, 130.4, 128.6, 125.7, 125.2, 116.3, 113.6, 62.5, 53.9, 45.5, 45.3, 36.2, 31.6, 30.6, 27.7, 24.0, 14.5.


**LRMS:** (*m/z*) 401 (M-H)<sup>+</sup>, 403 (M+H)<sup>+</sup>, 425 (M+Na)<sup>+</sup>.

**MIC (*E. coli*):** 6.25 μM

**MIC (*S. capitis*):** 1.6 μM

## Example 9

*N*-[1(*S*)-(Benzofuran-2-carbonyl)-2,2-dimethylpropyl]-3'-cyclopentyl-2'(*R*)-[(*N*-formyl-*N*-hydroxyamino)-methyl]-propionamide




The title compound was prepared as described in **Scheme 3** and the **General Protocol (GP)**. Experimental details and relevant spectroscopic data are described below.



Reagents and conditions:

**Step A:** 2-lithiobenzofuran, THF -70 to -40°C

## Scheme 3



Reagents and conditions:

**Step GPA:** HCl / dioxan

**Step GPB:** EDC, HOBT and either 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid or 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-3-cyclopentyl-propionic acid in DMF

**Step GPC:** H<sub>2</sub> / Pd/C 10 wt%

**General Protocol (GP)**

**STEP A:** [1(S)-(Benzofuran-2-carbonyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester.

A solution of benzofuran (4.24 g, 36 mmol) in dry THF (20 mL) under argon was cooled to  $-78^{\circ}\text{C}$ . *n*-Butyllithium (22.5 mL, 36 mmol, 1.6M in hexanes) was added dropwise such that an internal temperature below  $-65^{\circ}\text{C}$  was maintained. The reaction was allowed to stir at  $-78^{\circ}\text{C}$  for two hours during which time 2-lithiobenzofuran precipitated as a white solid. The suspension of 2-lithiobenzofuran was added dropwise to a cooled ( $-78^{\circ}\text{C}$ ) solution of [1(S)-(Methoxy-methyl-carbamoyl)-2,2-dimethylpropyl]-carbamic acid-*tert*-butyl ester (3.34 g, 12 mmol) in THF (30 mL) *via* cannula such that an internal temperature of below  $-65^{\circ}\text{C}$  was maintained. The yellow solution was allowed to stir at  $-70^{\circ}\text{C}$  for 30 minutes and  $-40^{\circ}\text{C}$  for a further 3 hours. The reaction was quenched by addition of saturated ammonium chloride solution (200 mL). The aqueous phase was extracted with ethyl acetate (3 x 150 mL) and the combined organic extracts washed with brine (2 x 200 mL), dried over magnesium sulphate, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography ( $\text{SiO}_2$ , 10 % ethyl acetate in hexanes) gave the title compound as a yellow solid (3.45 g, 87 %).

$^1\text{H NMR}$  ( $\text{CDCl}_3$ ):  $\delta/\text{ppm}$  7.70 (2H, m), 7.62 (1H, m), 7.50 (1H, m), 7.31 (1H, m), 5.41 (1H, bd,  $J = 9.1$  Hz), 5.07 (1H, bd,  $J = 9.7$  Hz), 1.43 (9H, s), 1.02 (9H, s).

LRMS: ( $m/z$ ) 354 ( $\text{M}+\text{Na}$ ) $^+$ .

**STEP GPA:** 2(S)-Amino-1-benzofuran-2-yl-3,3-dimethylbutan-1-one.

[1(S)-(Benzofuran-2-carbonyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester (700 mg, 2.11 mmol) was dissolved in 1,4-dioxan saturated with HCl (10 mL) and the yellow solution allowed to stir at room temperature for 18 hours.

The reaction was added to saturated sodium bicarbonate solution (50 mL) (CAUTION!) and the pH adjusted to 10 with the addition of 1M sodium carbonate.

The aqueous mixture was extracted with ethyl acetate (3 x 100mL) and the combined organic extracts washed with brine (3 x 100 mL), dried over magnesium sulphate, filtered and the solvents removed *in vacuo* to give the title compound as a pale yellow solid (480 mg, 98 %).

Alternatively the  $\alpha$ -aminoketones were isolated as hydrochloride salts by removal of the HCl / 1,4-dioxan *in vacuo*.

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm 7.72 (1H, d, J = 7.8 Hz), 7.55 (2H, m), 7.48 (1H, m), 7.32 (1H, t, J = 7.5 Hz), 4.12 (1H, s), 1.65 (2H, bs), 1.02 (9H, s).

**STEP GPB:** *N*-[1(*S*)-(Benzofuran-2-carbonyl)-2,2-dimethylpropyl]-3'-cyclopentyl-2'(*R*)-[(*N*-benzyloxy-*N*-formylamino)-methyl]-propionamide.

To a solution of 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-3-cyclopentyl-propionic acid (247 mg, 0.81 mmol) in DMF (5 mL) at 0 °C was added EDC (157 mg, 0.81 mmol) and HOBT (22 mg, 20 mol %) and the mixture allowed to stir at 0 °C for 45 minutes. A solution of 2(*S*)-Amino-1-benzofuran-2-yl-3,3-dimethylbutan-1-one (149 mg, 0.65 mmol) in DMF (2 mL) was added and the reaction allowed to stir at ambient temperature for 18 hours. The solvent was removed *in vacuo* and the residue dissolved in ethyl acetate (50 mL) and ammonium chloride solution (50 mL). The organic layer was separated and the aqueous phase extracted with ethyl acetate (2 x 50 mL). The combined organic extracts were dried over magnesium sulphate, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 2:1 hexanes : ethyl acetate) gave the title compound as a clear oil (196 mg, 58 %).

Alternatively where the hydrochloride salts of the  $\alpha$ -aminoketones were used the free amine was generated by the addition of 2 equivalents of TEA to the DMF solution of the  $\alpha$ -aminoketone prior to addition to the HOBT ester.

**<sup>1</sup>H NMR (MeOD):**  $\delta$ /ppm 8.42 – 8.25 (1H, bm), 8.13 (0.5H, m), 7.90 (0.5H, m), 7.78 (1H, s), 7.60 (1H, m), 7.59 – 7.30 (7H, m), 5.25 (1H, m), 4.90 (2H, m), 3.72 (1.5H, m), 3.33 (0.5H, m), 2.95 (1H, m), 1.75 (1H, m), 1.65 – 1.10 (8H, m), 0.98 (9H, s), 0.94 (2H, m).

**LRMS:** (*m/z*) 541 (M+Na)<sup>+</sup>

**STEP GPC:** *N*-[1(*S*)-(Benzofuran-2-carbonyl)-2,2-dimethylpropyl]-3'-cyclopentyl-2'(*R*)-[(*N*-formyl-*N*-hydroxyamino)-methyl]-propionamide.

*N*-[1(*S*)-(Benzofuran-2-carbonyl)-2,2-dimethylpropyl]-3'-cyclopentyl-2'(*R*)-[(*N*-benzyloxy-*N*-formylamino)-methyl]-propionamide (197 mg, 0.38 mmol) was dissolved in methanol (15 mL) under argon and a slurry of palladium on charcoal (20 mg) in ethyl acetate (1 mL) was added and hydrogen bubbled through the reaction for 10 minutes. An atmosphere of 1 bar of hydrogen was maintained via balloon and the reaction allowed to stir at ambient temperature for 1 hour. The reaction was filtered through glass wool and the solvents removed *in vacuo*. The residue was purified by reverse phase preparative HPLC to give the title compound as a white crystalline solid (84 mg, 52 %).

**<sup>1</sup>H NMR (MeOD):**  $\delta$ /ppm 8.28 (0.4H, s), 7.83 (2.6H, m), 7.65-7.45 (2H, m), 7.35 (1H, t, *J*=7.0Hz), 5.36 (1H, m), 3.77 (1H, dd, *J*=14.1, 9.5Hz), 3.61 (0.3H, dd, *J*=14.0, 5.4Hz), 3.43 (0.7H, dd, *J*=14.0, 4.6Hz), 3.10 (0.7H, m), 2.93 (0.3H, m), 1.82-1.15 (9H, m), 1.05 (9H, s), 1.10-0.80 (2H, m).

**<sup>13</sup>C NMR (MeOD):**  $\delta$ /ppm 191.5, 176.6, 157.8, 154.3, 130.4, 129.6, 125.7, 125.2, 116.3, 113.6, 62.5, 45.4, 39.4, 38.1, 36.3, 34.4, 33.8, 27.7, 26.3, 26.2.

**LRMS:** (*m/z*) 451 (M+Na)<sup>+</sup>.


**MIC (*E. coli*):** 6.25  $\mu$ M

**MIC (*S. capitis*):** 0.7  $\mu$ M




## Example 10

2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid pentafluorophenyl ester



The title compound was prepared as described in **Scheme 4**. The experimental details of the synthesis and relevant spectroscopic data are described below.



**Reagents and conditions:**

**Step A:** Formyl acetic anhydride, TEA, THF, 0 °C, 1 hour then RT, 1 hour;

**Step B:** Pentafluorophenol, EDC, EtOAc, 0 °C to RT, O/N.

**Scheme 4**

**Step A:** 2(*R*)-(N-Benzyloxy-*N*-formylamino)-methyl)-hexanoic acid

To a solution of 2(*R*)-(N-benzyloxyamino-methyl)-hexanoic acid (30.6 g, 0.12 mol) in dry THF (300 ml) was added formic acetic anhydride (26.8 ml, 0.31 mol) at 0 °C.

Triethylamine (18.5 ml, 0.13 mol) was added and the reaction mixture was stirred for 1 hour at 0 °C and 60 hours at room temperature. The solvent was removed *in vacuo* to yield the title compound as a yellow oil (33.6 g, 99%) which was used in **Step B** without further purification.

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm (rotamers) 8.14 (0.7H, br s), 8.00 (0.3H, br s), 7.50-7.25 (5H, m), 5.07-4.70 (2H, m), 3.95-3.52 (2H, m), 2.78 (1H, br s), 1.72-1.20 (6H, m) and 0.89 (3H, br s).

LRMS: (*m/z*) 280 (M+H)<sup>+</sup>.

**Step B:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid pentafluorophenyl ester


To a solution of 2(*R*)-(benzyloxy-formylamino)-methyl)-hexanoic acid (33.6 g, 0.12 mol) in dry THF (500 ml) was added pentafluorophenol (44.3 g, 0.24 mol), EDC (27.7 g, 0.14 mol) and hydroxybenzotriazole (16.2 g, 0.12 mol). The reaction was stirred overnight at room temperature. The solvent was removed *in vacuo* and the residue was dissolved in ethyl acetate, washed with 1 M Na<sub>2</sub>CO<sub>3</sub> (3 × 500 ml) and water (1 × 500 ml) and dried over MgSO<sub>4</sub> and concentrated *in vacuo* to yield a yellow oil (60 g). The product was purified by flash chromatography (SiO<sub>2</sub>, 5:1, hex:EtOAc to 1:2 hex:EtOAc) to yield a clear oil (42.0 g, 79%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm (rotamers) 8.15 (0.7H, br s), 8.01 (0.3H, br s), 7.60-7.21 (5H, m), 5.00-4.70 (2H, m), 4.04-3.72 (2H, m), 3.09 (1H, br s), 1.85-1.57 (2H, m), 1.50-1.26 (4H, m), 1.00-0.82 (3H, m).


LRMS: (*m/z*) 468 (M+Na), 446 (M+H)<sup>+</sup>.

## Example 11

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid [1'-(2,4-dimethoxybenzoyl)-2',2'-dimethylpropyl]-amide



The title compound was prepared as described in **Scheme 5**. The experimental details of the synthesis and relevant spectroscopic data are described below.



Reagents and conditions:

Step A: 4-lithio-3-methoxyanisole, THF -70°C

Step B: HCl / dioxan

Step C: 2-(*R*)-[(*N*-benzyloxy-*N*-formylamino)methyl]-hexanoic acid pentafluorophenyl ester, TEA and DCM.

Step D: H<sub>2</sub> / Pd / C

**Scheme 5**

**STEP A:** [1(S)-(2,4-Dimethoxy-benzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester.

To a solution of 1-bromo-2,4-dimethoxybenzene (482 mg, 2.22 mmol) in dry THF (10 mL) at  $-78^{\circ}\text{C}$  under argon was added dropwise *t*-BuLi (2.96 mL, 4.44 mmol, 1.5M in pentanes). After addition was complete the reaction mixture was allowed to stir at  $-75^{\circ}\text{C}$  for one hour. Meanwhile a solution of 2-*tert*-Butoxycarbonylamino-3,3-dimethyl-butyric acid pentafluorophenyl ester (Example 14) (400 mg, 1.01 mmol) in dry THF (5 mL) was prepared at  $-75^{\circ}\text{C}$  under argon. The solution of 4-lithio-3-methoxyanisole was then added dropwise via cannula such that an internal temperature of less than  $-70^{\circ}\text{C}$  was maintained. The reaction was allowed to stir at  $-70^{\circ}\text{C}$  for 4 and a half hours before saturated ammonium chloride solution (30 mL) was added. The aqueous phase was extracted with ethyl acetate (3 x 30 mL) and the combined organic extracts washed with 1M sodium carbonate solution (3 x 100 mL) and brine (2 x 100 mL). The ethyl acetate phase was dried over magnesium sulphate, filtered and the solvents removed *in vacuo* to give a clear oil. Purification by column chromatography ( $\text{SiO}_2$ , 6:1 hexanes : ethyl acetate) gave the title compound as a clear oil (239 mg, 68 %).

$^1\text{H NMR}$  ( $\text{CDCl}_3$ ):  $\delta/\text{ppm}$  7.70 (1H, d,  $J = 8.6$  Hz), 6.52 (1H, dd,  $J = 8.7, 2.2$  Hz), 6.42 (1H, m), 5.50 (1H, m), 5.40 (1H, d,  $J = 9.7$  Hz), 3.91 (3H, s), 3.86 (3H, s), 1.45 (9H, s), 0.88 (9H, s).

LRMS: ( $m/z$ ) 374 ( $\text{M}+\text{Na}^+$ )

**STEP B:** 2(S)-Amino-1-(2,4-dimethoxy-phenyl)-3,3-dimethyl-butan-1-one.HCl

To [1(S)-(2,4-Dimethoxy-benzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester (239 mg) at  $0^{\circ}\text{C}$  was added 1-4 dioxan saturated with HCl (3 mL). The reaction solution was allowed to stir for 18 hours before the solvents were removed *in vacuo* and the residue azeotroped with DCM (3 x 10 mL) to give the title compound as a yellow solid (224 mg).

**<sup>1</sup>H NMR (MeOD):**  $\delta$ /ppm 7.74 (1H, d,  $J = 9.3$  Hz), 6.89 (2H, m), 5.12 (1H, s), 3.97 (3H, s), 3.89 (3H, s), 0.99 (9H, s).

**LRMS:** ( $m/z$ ) 252 (M+H)<sup>+</sup>

**STEP C:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)methyl]-hexanoic acid [1'(*S*)-(2,4-dimethoxy-benzoyl)-2',2'-dimethylpropyl]-amide

To a solution of 2-{2-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoylamino}-3,3-dimethyl-butyric acid pentafluorophenyl ester (303 mg, 0.681 mmol) in DCM (3 mL) at 0 °C was added triethylamine (131 mg, 1.29 mmol) followed by a solution of 2(*S*)-Amino-1-(2,4-dimethoxy-phenyl)-3,3-dimethyl-butan-1-one.HCl in DCM (2 mL). The reaction was allowed to warm to ambient temperature and stirred for 18 hours. Saturated ammonium chloride solution (50 mL) was added and the aqueous phase extracted with ethyl acetate (3 x 30 mL). The organic phases were washed with water (100 mL), 1M sodium carbonate solution (3 x 100 mL) and brine (100 mL), dried over magnesium sulphate, filtered and the solvents removed *in vacuo* to give a crude brown oil. Purification by column chromatography (SiO<sub>2</sub>, 1:1 hexanes : ethyl acetate) gave the title product as a clear oil (153 mg 44 %).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):**  $\delta$ /ppm 8.13 (0.5, bs), 7.91 (0.5, bs), 7.69 (1H, bd,  $J = 8.5$  Hz), 7.60 – 7.30 (5H, m), 6.54 (1H, dd,  $J = 8.7, 2.3$  Hz), 6.40 (2H, m), 5.72 (1H, m), 5.00 and 4.80 (2H, m), 3.90 (3H, s), 3.85 (3H, s), 3.70 (1H, m), 3.10 (1H, m), 2.58 (1H, m), 1.55 – 1.10 (6H, m), 1.00 – 0.76 (12H, m).

**LRMS:** ( $m/z$ ) 513 (M+H)<sup>+</sup>, 535 (M+Na)<sup>+</sup>.

**STEP D:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)methyl]-hexanoic acid [1'(*S*)-(2,4-dimethoxy-benzoyl)-2',2'-dimethylpropyl]-amide

2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)methyl]-hexanoic acid [1'(*S*)-(2,4-dimethoxy-benzoyl)-2',2'-dimethylpropyl]-amide (154 mg, 0.30 mmol) was dissolved in methanol

(10 mL) under argon and a slurry of palladium on charcoal (15 mg) in ethyl acetate (1 mL) was added and hydrogen gas bubbled through the reaction solution for 10 minutes. The reaction was allowed to stir under a hydrogen atmosphere (balloon) for one hour. The reaction was filtered through glass wool and the solvents removed *in vacuo* to give the title compound as a white crystalline solid (99 mg, 78 %).

**<sup>1</sup>H NMR (MeOD):**  $\delta$ /ppm 8.27 (0.4H, s), 8.19 (0.6H, bd, J=9.0Hz), 8.10 (0.4H, bd, J=9.0Hz), 7.85 (0.6H, s), 7.63 (1H, m), 6.62 (2H, m), 5.72 (1H, m), 3.90 (3H, s), 3.86 (3H, s), 3.85 (1H, m), 3.61 (0.4H, m), 3.41 (0.6H, dd, J=14.2, 4.6Hz), 3.05 (0.6H, m), 2.90 (0.4H, m), 1.70 - 1.10 (6H, m), 0.90 (9H, bs), 0.82 (3H, m).

**<sup>13</sup>C NMR (MeOD):**  $\delta$ /ppm 176.3, 175.9, 166.8, 162.3, 133.8, 123.7, 107.3, 99.9, 64.9, 64.8, 56.6, 53.4, 46.2, 45.9, 36.2, 31.6, 30.6, 27.9, 24.0, 14.6.


**LRMS:** (*m/z*) 421 (M-H)<sup>+</sup>, 423 (M+H)<sup>+</sup>, 445 (M+Na)<sup>+</sup>.

**MIC (*E. coli*):** 25  $\mu$ M


**MIC (*S. capitis*):** 25  $\mu$ M

## Example 12

2(R)-[*(N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid [1'(S)-(4-hydroxy-benzoyl)-2',2'-dimethylpropyl]-amide



The title compound was prepared as described in **Scheme 6**. The experimental details of the synthesis and relevant spectroscopic data are described below.



Reagents and conditions:

Step A: Boron tribromide

Step B: EDC, HOBT and 2(R)-[*(N*-Benzyloxy-*N*-formylamino)-methyl]hexanoic acid, DMF

Step C: H<sub>2</sub> / Pd/C 10 wt%

**Scheme 6**

**STEP A: 2(S)-Amino-1-(4-hydroxy-phenyl)-3,3-dimethyl-butan-1-one**

To a solution of (1(S)-(4-Methoxybenzoyl)-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester (191 mg, 0.86 mmol) in 1,2-dichloroethane (15 mL) at 0 °C was added dropwise BBr<sub>3</sub> (3 mL, 3 mmol, 1.0 M in DCM). The brown solution was allowed to warm to room temperature and stirred for 18 hours. The reaction mixture was again cooled to 0 °C and 1M HCl in methanol (15 mL) added cautiously. The yellow solution was allowed to warm to ambient temperature and stirred for two hours before the solvents were removed *in vacuo* to give a yellow solid. The residue was dissolved in 1M HCl (30 mL) and washed with toluene (15 mL) and diethyl ether (15 mL). The pH of the aqueous phase was adjusted to 8 with the addition of solid sodium bicarbonate. The aqueous phase was then extracted with ethyl acetate (3 x 70 mL), dried over magnesium sulphate and filtered. The solvents were removed *in vacuo* to give the title compound as a pinkish solid (148 mg, 83 %).

<sup>1</sup>H NMR (MeOD): δ/ppm 7.88 (2H, d, J = 8.8 Hz), 6.83 (2H, d, J = 8.9 Hz), 4.32 (1H, s), 0.93 (9H, s).

LRMS: (m/z) 206 (M-H)<sup>-</sup>, 208 (M+H)<sup>+</sup>

**STEP B: 2(R)-[(N-Benzyloxy-N-formylamino)-methyl]-hexanoic acid [1'(S)-(4-hydroxy-benzoyl)-2,2-dimethylpropyl]-amide**

To a solution of 2(R)-[(N-Benzyloxy-N-formylamino)-methyl]-hexanoic acid (229 mg, 0.82 mmol) in DMF (10 mL) at 0 °C was added EDC (159 mg, 0.82 mmol) and HOBt (26 mg, 20 mol %) and the mixture allowed to stir at 0 °C for 30 minutes. A solution of 2(S)-Amino-1-(4-hydroxy-phenyl)-3,3-dimethyl-butan-1-one (140 mg, 0.68 mmol) in DMF (5 mL) was added and the reaction allowed to stir at ambient temperature for 18 hours. The solvent was removed *in vacuo* and the residue dissolved in ethyl acetate (30 mL) and water (70 mL). The organic layer was separated and the aqueous phase extracted with ethyl acetate (2 x 50 mL). The combined organic extracts were dried over magnesium sulphate, filtered and the solvents removed *in*



*vacuo* to give a crude off white foam. Purification by column chromatography (SiO<sub>2</sub>, 1:1 hexanes : ethyl acetate) gave the title compound as a white sticky foam (151 mg, 47 %).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 8.93 (1H, bs), 8.15 (0.5H, bs), 7.95 (0.5H, bs), 7.84 (2H, d, J = 8.8 Hz), 7.38 (5H, m), 6.88 (2H, d, J = 8.7 Hz), 6.48 (1H, bd, J = 9.1 Hz), 5.43 (1H, m), 4.90 (2H, bm), 3.75 (1.5H, bm), 3.12 (0.5H, bm), 2.60 (1H, m), 1.70 – 1.05 (6H, m), 0.93 (9H, s).

**LRMS:** (*m/z*) 491 (M+Na)<sup>+</sup>

**STEP C:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid [1'(*S*)-(4-hydroxy-benzoyl)-2',2'-dimethylpropyl]-amide

2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)methyl]-hexanoic acid [1'(*S*)-(4-hydroxy-benzoyl)-2',2'-dimethylpropyl]-amide (151 mg, 0.32 mmol) was dissolved in methanol (15 mL) under argon and a slurry of palladium on charcoal (20 mg) in ethyl acetate (1 mL) was added and hydrogen gas bubbled through the reaction solution for 10 minutes. The reaction was allowed to stir under a hydrogen atmosphere (balloon) for one hour. The reaction was filtered through glass wool and the solvents removed *in vacuo* to give the title compound as a white crystalline solid (119 mg, 98 %). Purification on the ASPEC preparative HPLC system gave the title product in >95 % purity (98 mg, 81 %).

**<sup>1</sup>H NMR (MeOD):** δ/ppm 8.27 (0.3H, s), 8.23 (0.7H, bd, J=9.4Hz), 8.11 (0.3H, bd, J=9.2Hz), 7.92 (2H, d, J=8.8Hz), 7.86 (0.7H, s), 6.85 (2H, d, J=8.8Hz), 5.48 (1H, m), 3.80 (1H, m), 3.61 (0.3H, dd, J=14.0, 5.7Hz), 3.41 (0.7H, dd, J=14.2, 4.6Hz), 3.02 (0.7H, m), 2.87 (0.3H, m), 1.60-1.30 (2H, m), 1.30-1.10 (4H, m), 0.96 (9H, s), 0.75 (3H, t, J=7.0Hz).

**<sup>13</sup>C NMR (MeOD):** δ/ppm 176.7, 176.2, 164.6, 132.7, 131.2, 116.8, 60.5, 60.4, 53.9, 45.8, 45.5, 36.3, 31.6, 30.6, 27.9, 24.0, 14.5.


**LRMS:** (*m/z*) 377 (M-H)<sup>-</sup>, 401 (M+Na)<sup>+</sup>.

MIC (*E. coli*): 6.25  $\mu$ M


MIC (*S. capitis*): 12  $\mu$ M

### Example 13

2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-3-cyclopentyl-propionic acid



The title compound was prepared as described in **Scheme 7**. The experimental details of the synthesis and relevant spectroscopic data are described below.



**Reagents and conditions:**

**Step A:** Formyl acetic anhydride, TEA, THF, 0 °C, 1 hour then RT, 1 hour;

**Scheme 7**

**Step A: 2(R)-[(N-Benzyloxy-N-formylamino)-methyl]-3-cyclopentyl-propionic acid**

2(R)-(Benzyloxyamino-methyl)-3-cyclopentyl-propionic acid was prepared as described above.


Formic acetic anhydride (27.78 g, 0.32 mol) followed by TEA (52.7 mL, 0.38mol) were added to a cooled (0 °C) solution of 2(R)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid in THF (180 mL). The reaction mixture was stirred for 1 hour at 0 °C and then at room temperature for 1 hour before the solvent was removed *in vacuo*. The residue was purified by column chromatography (SiO<sub>2</sub>, 25-100% EtOAc in hexanes) to give the title compound as a colourless oil (36.10 g, 94%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 10.50-10.15 (1H, m), 8.23-7.87 (1H, m), 7.50-7.10 (5H, m), 5.13-4.67 (2H, m), 3.97-3.54 (2H, m), 2.95-2.67 (1H, m), 2.13-0.93 (11H, m).


LRMS: (*m/z*) 328 (M + Na)<sup>+</sup>; 304 (M - H)<sup>-</sup>.

**Example 14**

2(S)-*tert*-Butoxycarbonylamino-3,3-dimethyl-butyric acid pentafluorophenyl ester



The title compound was prepared as described in Scheme 8. The experimental details of the synthesis and relevant spectroscopic data are described below.



Reagents and conditions:

A. Pentafluorophenol, NMM, EDC, EtOAc, 0 °C to RT, O/N.

### Scheme 8

**Step A:** 2(S)-tert-Butoxycarbonylamino-3,3-dimethyl-butyric acid pentafluorophenyl ester

2(S)-tert-Butoxycarbonylamino-3,3-dimethyl-butyric acid was prepared from L-tert-leucine according to **Step A** of **Scheme 1**.


Pentafluorophenol (8.75 g, 47.5 mmol) dissolved in EtOAc (5 mL) was added to a solution of 2(S)-tert-butoxycarbonylamino-3,3-dimethyl-butyric acid (10 g, 43 mmol) in EtOAc (35 mL) and the mixture cooled to 0 °C. NMM (4.75 mL, 43 mmol) followed by EDC (9.12 g, 47.6 mmol) were added to this cooled solution and the reaction mixture was then allowed to warm to room temperature and stirred overnight. The reaction mixture was then diluted with EtOAc (200 mL) and washed with 1M HCl (2 × 200 mL), 1M Na<sub>2</sub>CO<sub>3</sub> (3 × 200 mL) and brine, dried over MgSO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by column chromatography (SiO<sub>2</sub>, 10% EtOAc in hexanes) to give the title compound as a white solid (12.03 g, 70%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 5.16–4.99 (1H, m), 4.41 (1H, d, J = 9.4), 1.47 (9H, s) and 1.11 (9H, s).

LRMS: ( $m/z$ ) 420 ( $M + Na$ )<sup>+</sup>.


### Example 15

2(*R*)-[*(N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-*tert*-butyl-2'-oxo-3'-pyridin-2-yl-propyl) amide



The title compound was prepared as described in **Scheme 9**. The experimental details of the synthesis and relevant spectroscopic data are described below.

#### Scheme 9



Reagents and conditions:

**Step A.** (i) *n*-BuLi, THF, -40 °C, 1 hour (ii) -78 °C, 2(*S*)-*tert*-butoxycarbonylamino-3,3-dimethyl-butyric acid pentafluorophenyl ester /THF;

**Step B.** HCl/Dioxane, 2 hours, RT;

**Step C.** (i) 2(*R*)-[(benzyloxy-formyl-amino)-methyl]-hexanoic acid, EDC, HOBT, DMF, RT, 1.5 hours (ii) 3(*S*)-amino-4,4-dimethyl-1-pyridin-2-yl-pentan-2-one hydrochloride, NMM, DMF, RT, over W/E;

**Step D.** H<sub>2</sub>, Pd/C, MeOH.

**STEP A:** (1(*S*)-*tert*-Butyl-2-oxo-3-pyridin-2-yl-propyl)-carbamic acid *tert*-butyl ester

The reaction was carried out under argon in flame dried apparatus. *n*-BuLi (1.6 M in hexanes, 11.3 mL, 18.1 mmol) was added dropwise to a cooled (-40 °C) solution of 2-picoline (1.64 mL, 16.6 mmol) in THF (30 mL). When addition was complete the solution was stirred at -40 °C for 1 hour before cooling to -78 °C. A solution of 2(*S*)-*tert*-butoxycarbonylamino-3,3-dimethyl-butyric acid pentafluorophenyl ester (3.00 g, 7.6 mmol) in THF (30 mL) was added *via* a cannula, maintaining the reaction temperature below -65 °C. After stirring for 1 hour at -78 °C saturated NH<sub>4</sub>Cl solution (10 mL) was added and the reaction mixture was allowed to warm to room temperature before it was partitioned between EtOAc (200 mL) and saturated NH<sub>4</sub>Cl solution (200 mL). The organic extract was washed with water (200 mL) and brine (200 mL), dried over MgSO<sub>4</sub> and concentrated *in vacuo*. The residue was purified by column chromatography (SiO<sub>2</sub>, 25% Et<sub>2</sub>O in hexanes) to give the title product as a bright yellow solid (1.01 g, 44%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.57-8.51 (0.5H, m), 8.10 (0.5H, d, J = 5.2 Hz), 7.69-7.49 (1H, m), 7.25-7.13 (1H, m), 6.95-6.84 (1H, m), 5.40-5.28 (1H, m), 5.28-5.14 (0.5H, m), 4.29 (0.5H, d, J = 9.25 Hz), 4.09 (1H, s), 3.95 (0.5H, d, J = 9.8 Hz), 1.44 (4.5H, s), 1.42 (4.5H, s), 1.02 (4.5H, s) and 1.01 (4.5H, s).

[<sup>1</sup>H NMR is complex due to the presence of E and Z enol tautomers.]

**LRMS:** (*m/z*) 329 (M + Na)<sup>+</sup>.

**STEP B:** (*E*)- or (*Z*)-3(*S*)-Amino-4,4-dimethyl-1-pyridin-2-yl-pent-1-en-2-ol hydrochloride

(1(*S*)-*tert*-Butyl-2-oxo-3-pyridin-2-yl-propyl)-carbamic acid *tert*-butyl ester (300 mg, 1.0 mmol), was dissolved in dioxane saturated with HCl (20 mL) and the reaction mixture stirred at room temperature for 2 hours. The mixture was then concentrated *in vacuo* to give the title compound as a yellow solid (0.37 g) which was used in **Step C** without further purification.

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.91 (1H, d, J = 5.7 Hz), 8.73-8.60 (1H, m), 8.20-8.04 (2H, m), 5.16-4.77 (1H, m), 4.45 (1H, s), 1.23 (9H, s).

**STEP C:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid (1'(*S*)-*tert*-butyl-2'-oxo-3'-pyridin-2-yl-propyl) amide

EDC (0.16 g, 0.83 mmol) followed by HOBt (0.11 g, 0.81 mmol) was added to a solution of 2(*R*)-(benzyloxy-formylamino)-methyl-hexanoic acid (0.18 g, 0.64 mmol) in DMF (3 mL) and the reaction mixture stirred for 1.5 hours at room temperature. A mixture of (*E*)- or (*Z*)-3(*S*)-amino-4,4-dimethyl-1-pyridin-2-yl-pent-1-en-2-ol hydrochloride (1.0 mmol) and NMM (0.23 mL, 2.1 mmol) in DMF (1.5 mL) was heated until a solution was obtained and added to the reaction mixture which was then stirred at room temperature for 72 hours. The solvent was removed by concentration *in vacuo* and the residue partitioned between ethyl acetate (30 mL) and water (15 mL). The organic phase was washed with saturated sodium bicarbonate solution (15 mL) and brine (15 mL), dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified by column chromatography (SiO<sub>2</sub>, 50% ethyl acetate in hexanes) to give the title compound as a bright yellow solid (0.20 g, 68%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.58-6.21 (11H, m), 5.41-3.55 (6.5H, m), 2.66-2.46 (1H, m) and 1.73-0.67 (18H, m).

[<sup>1</sup>H NMR is complex due to the presence of E and Z enol tautomers.]

LRMS: (m/z) 468 (M + H)<sup>+</sup>; 466 (M – H)<sup>+</sup>.

**STEP D:** 2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(1'(S)-*tert*-butyl-2'-oxo-3'-pyridin-2-yl-propyl) amide

To a solution of 2(R)-[(N-benzyloxy-N-formylamino)-methyl]-hexanoic acid (1'(S)-*tert*-butyl-2'-oxo-3'-pyridin-2-yl-propyl) amide (0.20 g, 0.44mmol) in methanol (40 mL) under an argon atmosphere was added a slurry of 10% Pd/C (20 mg) in ethyl acetate (1 mL). Hydrogen gas was bubbled through the reaction mixture for 5 minutes before leaving the reaction to stir under 1 atmosphere of hydrogen for 10 hours. The reaction mixture was then flushed with argon and filtered through a pad of glass microfibre filters. The filtrate was concentrated *in vacuo* to give the title compound as a bright yellow glass (0.15 g, 92%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.73-6.65 (6H, m), 5.40-3.94 (2.5H, m), 3.94-3.24 (2H, m), 2.97-2.53 (1H, m), 1.73-1.10 (6H, m), [1.02, 1.00, 0.98 (9H, 3 × s)] and 0.89-0.63 (3H, m).

[<sup>1</sup>H NMR is complex due to the presence of E and Z enol tautomers.]


LRMS: (m/z) 378 (M + H)<sup>+</sup>; 376 (M – H)<sup>+</sup>.

MIC (*S. capitis*): 100 μM




**Example 16**

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid [1'(*S*)-(4-methanesulfonyl-benzoyl)-2',2'-dimethyl-propyl]amide



The title compound was prepared as described in **Scheme 10**. The experimental details of the synthesis and relevant spectroscopic data are described below.

**Reagents and conditions:**

**Step A:** MeSO<sub>2</sub>Na, DMSO, 130 °C, O/N;

**Step B:** HCl/Dioxane, O/N, RT;

**Step C:** (i) 2(*R*)-[(Benzyloxy-formylamino)-methyl]-hexanoic acid, EDC, HOBT, DMF, RT, 1.5 hours (ii) 2(*R*)- and 2(*S*)-amino-1-(4-methanesulfonyl-phenyl)-3,3-dimethyl-butan-1-one hydrochloride, NMM, DMF, RT, O/N;

**Step D:** (i) H<sub>2</sub>, Pd/C, MeOH, 3.5 hours, (ii) Prep. HPLC.

### Scheme 10

**STEP A:** Racemic mixture of 1(*R*)- and 1(*S*)-(4-Methanesulfonyl-benzoyl)-2,2-dimethyl-propyl-carbamic acid-*tert*-butyl ester

1(*S*)-(4-fluorobenzoyl)-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester was prepared as described in steps A to C in Scheme 1 (version 2).

Sodium methane sulphinat (132 mg, 1.3 mmol) was added to a solution of 1(*S*)-(4-fluorobenzoyl)-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester (200 mg, 0.6 mmol) in DMSO (4 mL) and the mixture stirred under argon at 130 °C for 17 hours. After cooling to room temperature the reaction mixture was partitioned between diethyl ether (100 mL) and water (100 mL), the organic extract washed with water (100 mL) and brine (100 mL), dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified by column chromatography (SiO<sub>2</sub>, 10% ethyl acetate in hexanes) to give the title compound as a white solid (0.22 g, 73%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.18 (2H, d, J = 8.4), 8.11-8.01 (2H, m), 5.42-5.27 (1H, m), 5.13 (1H, d, J = 9.6), 3.08 (3H, s), 1.43 (9H, s) and 0.94 (9H, s).

LRMS: (*m/z*) 392 (M + Na)<sup>+</sup>.

**STEP B:** 2(*R*)- and 2(*S*)-Amino-1-(4-methanesulfonyl-phenyl)-3,3-dimethyl-butan-1-one hydrochloride

1(*R*)- and 1(*S*)-(4-Methanesulfonyl-benzoyl)-2,2-dimethyl-propyl-carbamic acid-*tert*-butyl ester (0.22 g, 0.6 mmol), was dissolved in dioxane saturated with HCl (20 mL)

and the reaction mixture stirred at room temperature for 17 hours. The mixture was then concentrated *in vacuo* to give the title compound as a yellow solid which was used in **Step C** without further purification.

**<sup>1</sup>H NMR (MeOD):**  $\delta$ /ppm 8.34–8.12 (4H, m), 5.12 (1H, s), 3.20 (3H, s) and 1.04 (9H, s).

**STEP C: (2*R*,1'*R*)- and (2*R*,1'*S*)-2-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid [1'-(4-methanesulfonyl-benzoyl)-2',2'-dimethylpropyl]amide**

EDC (98 mg, 0.51 mmol) followed by HOBt (70 mg, 0.52 mmol) were added to a solution of 2(*R*)-[(benzyloxy-formylamino)-methyl]-hexanoic acid (0.11 g, 0.39 mmol) in DMF (2 mL) and the reaction mixture stirred for an hour at room temperature. 2(*R*)- and 2(*S*)-amino-1-(4-methanesulfonyl-phenyl)-3,3-dimethyl-butan-1-one hydrochloride (0.6 mmol) and NMM (0.13 mL, 1.2 mmol) in DMF (1.0 mL) were heated until a solution was obtained, added to the reaction mixture and stirred overnight. The solvent was removed by concentration *in vacuo* and the residue partitioned between ethyl acetate (40 mL) and 1M HCl (40 mL). The organic phase was washed with saturated sodium bicarbonate solution (40 mL) and brine (40 mL), dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified by column chromatography (SiO<sub>2</sub>, 1.5% methanol in DCM) to give the title compound as a white solid (140 mg, 67%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):**  $\delta$ /ppm 8.30–7.92 (5H, m), 7.51–7.10 (5H, m), 6.50–6.17 (1H, m), 5.66–5.49 (0.4H, m), 5.49–5.33 (0.6H, m), 5.10–4.45 (2H, m), 3.93–3.53 (2H, m), 3.21–2.97 (3H, m), 2.68–2.49 (1H, m), 1.61–1.03 (6H, m) and 1.04–0.73 (12H, m).

**LRMS: (*m/z*)** 553 (M + Na)<sup>+</sup>, 531 (M + H)<sup>+</sup>.

**STEP D: (2*R*,1'*R*)- and (2*R*,1'*S*)-2-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid [1'-(4-methanesulfonyl-benzoyl)-2',2'-dimethyl-propyl]amide**

To a solution of (2*R*,1'*R*)- and (2*R*,1'*S*)-2-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid [1'-(4-methanesulfonyl-benzoyl)-2',2'-dimethylpropyl]amide (140 mg, 0.26 mmol) in methanol (30 mL) under an argon atmosphere was added a slurry of 10% Pd/C (14 mg) in ethyl acetate (1 mL). Hydrogen gas was bubbled through the reaction mixture for 5 minutes before leaving the reaction to stir under 1 atmosphere of hydrogen for 3.5 hours. The reaction mixture was then flushed with argon and filtered through a pad of glass microfibre filters. The filtrate was concentrated *in vacuo* and the residue purified by reverse phase HPLC to give the (2*R*, 1'*S*) isomer as a white solid (32 mg, 28%).

<sup>1</sup>H NMR (MeOD): δ/ppm 8.24 (2H, d, J = 8.5), 8.10 (2H, d, J = 8.5), 7.95-7.56 (1H, m), 5.49-5.36 (1H, m), 3.86-3.36 (2H, m), 3.17 (3H, s), 3.15-2.80 (1H, m), 1.61-1.04 (6H, m), 1.04-0.90 (9H, m) and 0.84-0.78 (3H, m).

<sup>13</sup>C NMR (MeOD): δ/ppm (rotamers) 201.2, 201.1, 176.9, 176.6, 175.6, 146.3, 144.3, 144.1, 130.8, 129.9, 129.4, 128.6, 76.2, 62.7, 62.2, 53.9, 53.5, 45.4, 45.2, 44.5, 42.0, 36.0, 35.8, 32.4, 31.6, 30.5, 30.0, 28.7, 28.4, 24.0 and 14.6.

LRMS: (*m/z*) 463 (M + Na)<sup>+</sup>, 441 (M + H)<sup>+</sup>; 439 (M - H)<sup>-</sup>.


HPLC: 82%.

MIC (*E. coli*): 6.25 μM

MIC (*S. capitis*): 25 μM

**Example 17**

3-Cyclopentyl-2(S)-[(*N*-formyl-*N*-hydroxyamino)-methyl]-*N*-[1'(*R*)-(4-methoxybenzoyl)-2',2'-dimethylpropyl]propionamide [Enantiomer of **Example 5**]



The title compound was prepared in an analogous manner to **Example 5** as described in **Scheme 1** utilising D-*t*-leucine in **Step A** and 2(*S*)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid in **Step E**. 2(*S*)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid was synthesised in an analogous manner to 2(*R*)-(benzyloxyamino-methyl)-3-cyclopentyl-propionic acid using (*R*)-(+)-4-benzyl-2-oxazolidinone as the chiral auxilliary.


Spectral data was consistent with **Example 5**.

MIC (*E. coli*): 100  $\mu$ M

MIC (*S. capitis*): 25  $\mu$ M

**Example 18**

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(furan-2-carbonyl)-2',2'-dimethylpropyl)amide



The title compound was prepared from furan in an analogous manner to that described in **Scheme 2**, using 2-lithiofuran in **Step C**. The relevant spectroscopic data are described below.

**STEP C:** 2(*R*)-(*N*-Benzyloxyaminomethyl)-hexanoic acid-(1'(*S*)-(furan-2-carbonyl)-2',2'-dimethylpropyl)amide.

Purification by column chromatography (SiO<sub>2</sub>, 5:1 *iso*-hexane:ethyl acetate) gave the product as a colourless oil (75 mg, 37%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.69-7.63 (1H, m), 7.46-7.30 (5H, m), 7.33 (1H, d, J = 3.5 Hz), 6.98 (1H, d, J = 9.6 Hz), 6.57 (1H, dd, J = 3.5, 1.7 Hz), 5.72 (1H, brs), 5.34 (1H, d, J = 9.6 Hz), 4.75 (2H, s), 3.22-3.01 (2H, m), 2.60-2.44 (1H, m), 1.69-1.50 (1H, m), 1.50-1.34 (1H, m), 1.34-1.12 (4H, m), 1.01 (9H, s), 1.84 (3H, t, J = 6.8 Hz).

LRMS: (*m/z*) 414.8 (M+H)<sup>+</sup>.

**STEP D:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)methyl]-hexanoic acid-(1'(*S*)-(furan-2-carbonyl)-2',2'-dimethylpropyl)amide.

Purification by column chromatography (SiO<sub>2</sub>, 4:1 to 2:1 *iso*-hexane:ethyl acetate) gave the product as a colourless oil (70 mg, 87%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 8.04 (0.6H, brs), 7.90 (0.4H, brs), 7.65 (1H, d, *J* = 1.7 Hz), 7.52-7.32 (5H, m), 7.32 (1H, d, *J* = 3.5 Hz), 6.28 (1H, dd, *J* = 3.5, 1.7 Hz), 6.29 (1H, d, *J* = 9.3 Hz), 5.27 (1H, d, *J* = 9.2 Hz), 5.10-4.90 (1H, m), 4.90-4.72 (1H, m), 3.87-3.58 (2H, m), 3.20-3.01 (1H, m), 2.74-2.50 (1H, m), 1.65-1.46 (2H, m), 1.46-1.06 (4H, m), 0.95 (9H, s), 0.80 (3H, t, *J* = 6.8 Hz).

**LRMS:** (*m/z*) 423.2 (*M*+H)<sup>+</sup>, 465.2 (*M*+Na)<sup>+</sup>.

**STEP E:** 2(*R*)-[(*N*-formyl-*N*-hydroxyamino)methyl]-hexanoic acid-(1'(*S*)-(furan-2-carbonyl)-2',2'-dimethylpropyl)amide.

The title compound was isolated as a white crystalline solid (50 mg, 84%).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):** δ/ppm 8.35 (0.6H, d, *J* = 8.8 Hz), 8.26 (0.2H, s), 8.24 (0.4H, d), 7.89-7.82 (1.8H, m), 7.46 (1H, d, *J* = 3.6 Hz), 6.67 (1H, dd, *J* = 3.6, 1.7 Hz), 5.28-5.20 (1H, m), 3.85-3.70 (1H, m), 3.61 (0.4H, dd, *J* = 14.0, 5.5 Hz), 3.41 (0.6H, dd, *J* = 14.2, 4.6 Hz), 3.12-2.98 (0.6H, m), 2.96-2.82 (0.4H, m), 1.57-1.36 (2H, m), 1.36-1.12 (4H, m), 1.00 (3.6H, s), 0.99 (5.4H, s), 0.80 (3H, t, *J* = 6.9 Hz).

**<sup>13</sup>C NMR (CD<sub>3</sub>OD):** δ/ppm 189.5, 176.7, 176.4, 154.4, 149.7, 120.9, 114.2, 62.2, 62.1, 53.9, 50.3, 45.6, 45.3, 36.3, 31.6, 30.6, 27.7, 24.0, 14.6.


**LRMS:** (*m/z*) 353.2 (*M*+H)<sup>+</sup>, 375.2 (*M*+Na)<sup>+</sup>, 351.2 (*M*-H)<sup>-</sup>.

**MIC (*E. coli*):** 12.5 μM

**MIC (*S. capitis*):** 12.5 μM

**Example 19**

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-(1-methyl-1*H*-pyrrole-2-carbonyl)-propyl)amide



The title compound was prepared from 1-methylpyrrole in an analogous manner to that described in **Scheme 5**, using 2-lithio-1-methylpyrrole in **Step A**. The relevant spectroscopic data are described below.

**STEP B:** [2,2-dimethyl-1(*S*)-(1-methyl-1*H*-pyrrole-2-carbonyl)-propyl]-carbamic acid *tert*-butyl ester.

Purification by column chromatography (SiO<sub>2</sub>, 18:1 *iso*-hexane:ethyl acetate) gave the product as an orange oil (0.18 g, 80%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 7.19-7.12 (1H, m), 6.90-6.84 (1H, m), 6.19-6.12 (1H, m), 5.48 (1H, d, *J* = 9.6 Hz), 4.82 (1H, d, *J* = 9.6 Hz), 3.94 (3H, s), 1.44 (9H, s), 0.98 (9H, s).

**LRMS:** (*m/z*) 295.2 (M+H)<sup>+</sup>.

**STEP C:** 2(*S*)-Amino-3,3-dimethyl-1-(1-methyl-1*H*-pyrrol-2-yl)-butan-1-one hydrochloride.



The title compound was isolated as a pale purple solid and used without further purification (0.13g, 100% crude).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 7.15-7.08 (1H, m), 6.93-6.87 (1H, m), 6.18-6.10 (1H, m), 4.61 (1H, brs), 3.93 (3H, s), 1.02 (9H, s).

LRMS: (m/z) 195.0 (M+H)<sup>+</sup>.

**STEP D:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-(1-methyl-1*H*-pyrrole-2-carbonyl)-propyl)amide.

Purification by column chromatography (SiO<sub>2</sub>, 4:1 to 1:1 *iso*-hexane:ethyl acetate) gave the product as a pale brown oil (0.16 g, 68%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.13 (0.6H, brs), 7.92 (0.4H, brs), 7.56-7.28 (5H, m), 7.21-7.12 (1H, m), 6.92-6.83 (1H, m), 6.31 (1H, d, J = 9.7 Hz), 6.25 (1H, dd, J = 3.8, 2.6 Hz), 5.17 (1H, d, J = 9.7 Hz), 5.08-4.91 (1H, m), 4.91-4.73 (1H, m), 3.92 (3H, s), 3.89-3.54 (2H, m), 3.19-3.02 (1H, m), 2.64-2.48 (1H, m), 1.64-1.32 (2H, m), 1.32-1.07 (4H, m), 0.96 (9H, s), 0.80 (3H, t, J = 7.0 Hz).

LRMS: (m/z) 456.2 (M+H)<sup>+</sup>, 478.2 (M+Na)<sup>+</sup>.

**STEP E:** 2(*R*)-[(*N*-formyl-*N*-hydroxyamino)methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-(1-methyl-1*H*-pyrrole-2-carbonyl)-propyl)amide.

The title compound was isolated as a pale brown crystalline solid (60 mg, 80%).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.26 (0.5H, brs), 8.18 (0.5H, d, J = 9.4 Hz), 8.07 (0.5H, brd, J = 10.1 Hz), 7.85 (0.5H, s), 7.19 (1H, d, J = 3.8 Hz), 7.10-7.00 (1H, m), 6.15 (1H, dd, J = 3.7, 2.4 Hz), 5.25-5.15 (1H, m), 3.90 (3H, s), 3.88-3.71 (1H, m), 3.70-3.52 (0.5H, m), (0.5H obs. by CD<sub>3</sub>OD peak), 3.11-2.92 (0.5H, m), 2.92-2.76 (0.5H, m), 1.63-1.31 (2H, m), 1.31-1.05 (4H, m), 0.99 (4.5H, s), 0.97 (4.5H, s), 0.76 (3H, t, J = 6.8 Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 190.5, 176.4, 176.1, 134.5, 132.7, 122.9, 109.7, 61.9, 61.8, 53.9, 50.2, 45.8, 45.6, 38.3, 36.4, 31.6, 30.6, 28.1, 24.0, 14.6.


LRMS: (*m/z*) 366.2 (M+H)<sup>+</sup>, 388.2 (M+Na)<sup>+</sup>, 363.8 (M-H)<sup>-</sup>.

MIC (*E. coli*): 12.5 μM


MIC (*S. capitis*): 12.5 μM

### Example 20

2(*R*)-[*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-aminobenzoyl)-2',2'-dimethylpropyl)amide



The title compound was prepared in as described in **Scheme 11**, using the **General Protocol (GP)** thereafter. The experimental details of the synthesis and relevant spectroscopic data are described below.



## Reagents and Conditions:

A.  $\text{NaN}_3$ , DMF

B. Benzyl chloroformate, pyridine, DCM

## Scheme 11

**STEP A:** [1(S)-(4-Aminobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester.

Sodium azide (1.26 g, 19.4 mmol) was added to a solution of [1(S)-(4-fluorobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester (0.20 g, 0.7 mmol) in DMF (15 mL), and the mixture heated at 120 °C for 40 hours. The solvent was removed *in vacuo* before addition of 10%  $\text{NaHCO}_3$  (aq.) (30 mL) to the residual solid. The aqueous phase was extracted with ethyl acetate (2 x 30 mL) and the organic phases washed with brine (1 x 30 mL), dried over  $\text{MgSO}_4$ , filtered and the solvent removed *in vacuo* to give a yellow solid. Purification by column chromatography ( $\text{SiO}_2$ , 4:1 *iso*-hexane:ethyl acetate) afforded the title compound as a yellow solid (67 mg, 36%).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$ /ppm 7.85 (2H, d,  $J$  = 8.5 Hz), 6.65 (2H, d,  $J$  = 8.6 Hz), 5.45 (1H, brd,  $J$  = 9.2 Hz), 5.08 (1H, d,  $J$  = 9.6 Hz), 4.17 (2H, brs), 1.42 (9H, s), 0.93 (9H, s).

LRMS: ( $m/z$ ) 329.2 ( $\text{M}+\text{Na}$ ) $^+$ .

**STEP B:** [4-(2(*S*)-*tert*-Butoxycarbonylamino-3,3-dimethylbutyryl)-phenyl]-carbamic acid benzyl ester.

A solution of [1(*S*)-(4-aminobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester (67 mg, 0.22 mmol) and triethylamine (95  $\mu$ L, 0.69 mmol) in DCM (3 mL) was cooled on an ice bath prior to addition of benzyl chloroformate (98  $\mu$ L, 0.69 mmol). The reaction was warmed to room temperature and left for 15 hours. Examination by T.l.c. suggested minimal conversion to product hence pyridine (0.10 mL, 1.2 mmol) was added and the reaction left for a further 2 hours. Solvent was removed *in vacuo* and the residue taken up in ethyl acetate (20 mL). The organic phase was washed with 1M HCl (1 x 20 mL), brine (1 x 20 mL), 1M Na<sub>2</sub>CO<sub>3</sub> (1 x 20 mL) and brine (1 x 20 mL, dried over MgSO<sub>4</sub>, filtered and the solvent removed *in vacuo*, giving a yellow solid. Purification by column chromatography (SiO<sub>2</sub>, 9:1 *iso*-hexane:ethyl acetate) afforded the title compound as a yellow solid (90 mg, 90%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):**  $\delta$ /ppm 7.77 (1H, d, J = 8.7 Hz), 7.50 (1H, d, J = 8.7 Hz), 7.42-7.32 (5H, m), 7.00 (1H brs), 5.43 (1H, brd, J = 9.5 Hz), 5.22 (2H, s), 5.12 (1H, d, J = 9.7 Hz), 1.42 (9H, s), 0.91 (9H, s).

**LRMS:** (*m/z*) 463.2 (M+Na)<sup>+</sup>, 439.2 (M-H)<sup>-</sup>.

**STEP GPA:** [4-(2(*S*)-amino-3,3-dimethylbutyryl)-phenyl]-carbamic acid benzyl ester hydrochloride.

The title compound was isolated as a foam and used without further purification (75 mg, 100% crude).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):**  $\delta$ /ppm 8.88 (0.5H, brs), 8.41-8.21 (3H, m), 7.59-7.50 (2.5H, m), 7.30-7.21 (2H, m, obs. by CDCl<sub>3</sub> peak), 7.432-7.31 (5H, m), 7.00 (1H brs), 4.86-4.68 (1H, m), 5.15 (2H, s), 1.04 (9H, s).

**LRMS:** (*m/z*) 341.2 (M+H)<sup>+</sup>, 363.2 (M+Na)<sup>+</sup>.

**STEP GPB:** [4-(2(*S*)-{2(*R*)-(N-Formyl-N-hydroxyamino)-methyl]-hexanoylamino)-3,3-dimethylbutyryl)-phenyl]-carbamic acid benzyl ester.

Purification by column chromatography (SiO<sub>2</sub>, 4:1 *iso*-hexane:ethyl acetate) afforded the title compound as a pale brown oil (50 mg, 42%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 8.13 (0.4H, brs), 8.04-7.80 (2.6H, m), 7.78-7.52 (1H, m), 7.57 (2H, d, J = 8.7), 7.45-7.28 (10H, m), 6.38 (1H, d, J = 9.3 Hz), 5.46 (1H, d, J = 9.3 Hz), 5.20 (2H, s), 5.08-4.86 (1H, m), 4.86-4.70 (1H, m), 3.89-3.54 (2H, m), 2.62-2.50 (1H, m), 1.62-1.27 (2H, m), 1.27-1.05 (4H, m), 0.88 (9H, s), 0.75 (3H, t, J = 6.9 Hz).

**LRMS:** (*m/z*) 602.2 (M+H)<sup>+</sup>, 624.2 (M+Na)<sup>+</sup>, 600.2 (M-H)<sup>-</sup>.

**STEP GPC:** 2(*R*)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-aminobenzoyl)-2',2'-dimethylpropyl)amide.

The title compound was isolated as a white crystalline solid (29.5 mg, 100%).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):** δ/ppm 8.27 (0.3H, s), 8.11 (0.7H, d, J = 9.3 Hz), 8.00 (0.3H, d, J = 9.4 Hz), 7.86 (0.7H, s), 7.80 (2H, d, J = 8.8 Hz), 6.65 (2H, d, J = 8.8 Hz), 5.44 (1H, d, J = 9.4 Hz), 3.8 (1H, dd; J = 14.2, 9.7 Hz), 3.61 (0.5H, dd, J = 13.8, 5.4 Hz), 3.41 (0.5H, dd, J = 14.2, 4.6 Hz), 3.09-2.93 (1H, m), 1.59-1.30 (2H, m), 1.30-1.03 (4H, m), 0.96 (4.5H, s), 0.95 (4.5H, s), 0.75 (3H, t, J = 6.9 Hz).

**<sup>13</sup>C NMR (CD<sub>3</sub>OD):** 199.2, 176.4, 176.1, 156.1, 132.8, 127.6, 114.7, 29.9, 53.9, 50.3, 45.9, 45.6, 36.4, 31.9, 30.6, 28.0, 24.0, 14.5.


**LRMS:** (*m/z*) 400.2 (M+Na)<sup>+</sup>.

**MIC (*E. coli*):** 6.25 μM


**MIC (*S. capitis*):** 12.5 μM

## Example 21

2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)-[4-(2,2,2-trifluoroacetyl-amino)-benzoyl]-propyl)amide



The title compound was prepared as described in **Scheme 12**, using the **General Procedure (GP)** thereafter. The experimental details of the synthesis and relevant spectroscopic data are described below.



## Reagents and Conditions:

A.  $\text{NaN}_3$ , DMF,  $120^\circ\text{C}$

B. i)  $\text{H}_2$ , 10% Pd-C, MeOH, ii) Trifluoroacetic anhydride, pyridine, DCM

## Scheme 12

**STEP A:** [1(S)-(4-Azidobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester.

Sodium azide (37.9 g, 0.58 mol) was added to a solution of [1(S)-(4-fluorobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester (6.0 g, 19.4 mmol) in DMF (400 mL), and the mixture heated at 120 °C for 24 hours. The solvent was removed *in vacuo* and the residual solid taken up in ethyl acetate (200 ml). The organic phase was washed with 1M Na<sub>2</sub>CO<sub>3</sub> (1 x 400 ml) and brine (1 x 400 mL), dried over MgSO<sub>4</sub>, filtered and the solvent removed *in vacuo* to give a yellow solid. Purification by column chromatography (SiO<sub>2</sub>, 6:1 *iso*-hexane:ethyl acetate) afforded the title compound as a yellow solid (1.96 g) and [1(S)-(4-Azidobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester as an orange solid (0.97 g) (47% combined yield).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.01 (2H, d, J = 8.7 Hz), 7.09 (2H, d, J = 8.7 Hz), 5.37 (1H, brd, J = 9.4 Hz), 5.10 (1H, d, J = 9.6 Hz), 1.43 (9H, s), 0.93 (9H, s).

LRMS: (*m/z*) 327.2 (M+H)<sup>+</sup>, 355.2 (M+Na)<sup>+</sup>.

**STEP B:** (2,2-dimethyl-1(S)-[4-(2,2,2-trifluoroacetyl-amino)-benzoyl]-propyl)-carbamic acid *tert*-butyl ester.

To a degassed solution of [1(S)-(4-azidobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester (1.0 g, 3.0 mmol) in methanol (10 mL) was added 10% palladium on charcoal (0.20 g). Hydrogen gas was bubbled through the suspension for 15 minutes when a balloon containing hydrogen was introduced to the reaction. After 15 hours, the catalyst was removed by filtration and washed liberally with methanol and DCM. The solvents were removed *in vacuo* to give a pale cream solid (1.0 g). The crude material was mixed with DCM (20 mL) and dissolution occurred on addition of triethylamine (0.63 mL, 3.16 mmol). The mixture was cooled to 0°C (ice bath) and trifluoroacetic anhydride (0.45 mL, 3.16 mmol) was added dropwise. After 2 hours, further triethylamine (0.3 mL, 2.2 mmol) and trifluoroacetic anhydride (0.30 mL, 2.11 mmol) were added and the reaction proceeded for a further 2 hours. The solvent was removed *in vacuo* and ethyl acetate (20 mL) added to the residue. The organic phase was washed with 10% citric acid (aq.) (1 X 40 mL) and brine (1 x 40 mL), dried over MgSO<sub>4</sub>, filtered and the solvent removed *in vacuo* to give a brown

oil. Purification by column chromatography (SiO<sub>2</sub>, 9:1 *iso*-hexane:ethyl acetate) afforded the title compound as a pale brown oil (0.63 g, 52%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.58 (1H, brs), 8.00 (2H, d, J = 8.6 Hz), 7.69 (2H, dd, J = 8.5, 1.9 Hz), 5.42 (1H, brd, J = 9.4 Hz), 5.10 (1H, d, J = 9.5 Hz), 1.43 (9H, s), 0.93 (9H, s).

LRMS: (*m/z*) 425.2 (M+Na)<sup>+</sup>.

**STEP GPA:** *N*-[4-(2(*S*)-Amino-3,3-dimethylbutyryl)-phenyl]-2,2,2-trifluoroacetamide.

The title compound was recovered as a pale brown solid and used without further purification (0.55 g, 100% crude).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.12 (2H, d, J = 8.8 Hz), 7.90 (2H, d, J = 8.8 Hz), 5.04 (1H, s), 1.05 (9H, s).

LRMS: (*m/z*) 303.2 (M+H)<sup>+</sup>, 325.2 (M+Na)<sup>+</sup>.

**STEP GPB:** 2(*R*)-[*N*-Benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(2,2,2-trifluoroacetyl-amino)-benzoyl]-propyl)-amide.

Purification by column chromatography (SiO<sub>2</sub>, 8:1 *iso*-hexane:ethyl acetate) afforded the title compound as a pale brown oil (0.4 g, 44%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 9.03 (1H, brs), 8.13 (0.6H, brs), 8.00 (2H, d, J = 8.7), 7.91 (0.4H, brs), 7.72 (2H, d, J = 8.8), 7.43-7.32 (5H, m), 6.36 (1H, d, J = 8.9 Hz), 5.42 (1H, d, J = 9.0 Hz), 5.08-4.90 (1H, m), 4.90-4.74 (1H, m), 3.95-3.75 (1H, m), 3.75-3.58 (1H, m), 2.68-2.55 (1H, m), 1.62-1.30 (2H, m), 1.30-1.05 (4H, m), 0.88 (9H, s), 0.78 (3H, t, J = 6.9 Hz).

LRMS: (*m/z*) 564.2 (M+H)<sup>+</sup>, 686.2 (M+Na)<sup>+</sup>.



**STEP GPC:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(2,2,2-trifluoroacetyl-amino)-benzoyl]-propyl)amide

Purification by reverse phase preparative HPLC yielded the title compound as a white glassy solid (20 mg, 50%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.27 (0.3H, s), 8.06 (2H, d, *J* = 8.7 Hz), 7.90-7.78 (2.7H, m), 5.46 (1H, s), 3.88-3.72 (1H, m), 3.61 (0.3H, dd, *J* = 14.2, 5.8 Hz), 3.42 (0.7H, dd, *J* = 14.1, 4.3 Hz), 3.10-2.96 (0.7H, m), 2.95-2.80 (0.3H, m), 1.59-1.30 (2H, m), 1.30-1.04 (4H, m), 0.99 (3H, s), 0.98 (6H, s) and 0.75 (3H, t, *J* = 6.9 Hz).

<sup>13</sup>C NMR (CDCl<sub>3</sub>): δ/ppm 200.6, 200.4, 177.0, 176.7, 176.4, 157.3 (q, *J* = 38 Hz), 142.9, 136.7, 131.2, 122.0, 117.7 (q, *J* = 288 Hz), 61.1, 45.6, 45.4, 36.1, 31.6, 30.5, 27.8, 24.0, 14.5.


LRMS: (*m/z*) 474.2 (M+H)<sup>+</sup>, 496.2 (M+Na)<sup>+</sup>, 472.2 (M-H)<sup>-</sup>.

MIC (*E. coli*): 6.25 μM


MIC (*S. capitis*): 6.25 μM

### Example 22

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-acetylaminobenzoyl)-2',2'-dimethylpropyl)amide



The title compound was prepared from 2(*R*)-[(*N*-benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(2,2,2-trifluoroacetyl-amino)-benzoyl]-propyl)-amide as described in **Scheme 13**. The experimental details of the synthesis and relevant spectroscopic data are described below.



**Reagents and Conditions:**

- A. 1M Na<sub>2</sub>CO<sub>3</sub> (aq.), MeOH
- B. Acetic anhydride, pyridine, DCM
- C. H<sub>2</sub>, 10% Pd-C, MeOH

**Scheme 13**

**STEP A:** 2(*R*)-[(*N*-Benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-[4-aminobenzoyl]-2',2'-dimethylpropyl)-amide.

To a solution of 2(*R*)-[(*N*-benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(2,2,2-trifluoroacetyl-amino)-benzoyl]-propyl)-amide (0.35 g, 0.62 mmol) in methanol (4 mL) was added 1M Na<sub>2</sub>CO<sub>3</sub>. The reaction was left at room temperature for 4 hours when the mixture was extracted with ethyl acetate (2 x 20

mL). The organic phases were washed with brine (1 x 20 mL) and dried over  $\text{MgSO}_4$ , filtered and the solvent removed *in vacuo*. The resulting foam was purified by column chromatography ( $\text{SiO}_2$ , 4:1 *iso*-hexane:ethyl acetate) afforded the title compound as a pale brown oil (0.23 g, 79%).

**$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):**  $\delta$ /ppm 9.03 (1H, brs), 8.13 (0.6H, brs), 7.91 (0.4H, brs), 7.85 (2H, d,  $J = 8.67$ ), 7.46-7.32 (5H, m), 6.64 (2H, d,  $J = 8.7$ ), 6.38 (1H, d,  $J = 9.3$  Hz), 5.43 (1H, d,  $J = 9.4$  Hz), 5.09-4.90 (1H, m), 4.90-4.74 (1H, m), 3.90 (2H, m), 2.62-2.48 (1H, m), 1.612-1.28 (2H, m), 1.28-1.03 (4H, m), 0.90 (9H, s), 0.76 (3H, t,  $J = 6.9$  Hz).  
**LRMS:** ( $m/z$ ) 468.2 ( $\text{M}+\text{H}$ ) $^+$ , 490.2 ( $\text{M}+\text{Na}$ ) $^+$ , 466.4 ( $\text{M}-\text{H}$ ) $^-$ .

**STEP B:** 2(*R*)-[(*N*-Benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-(4-acetylaminobenzoyl)-2',2'-dimethylpropyl)-amide.

A solution of 2(*R*)-[(*N*-benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-(4-aminobenzoyl)-2',2'-dimethylpropyl)-amide (0.1 g, 0.21 mmol) in DCM (2 mL) was cooled to 0°C (ice bath) before addition of pyridine (34.6  $\mu\text{L}$ , 0.43 mmol) and acetyl chloride (30.5  $\mu\text{L}$ , 0.43 mmol). The reaction was warmed to room temperature and left for 15 hours. The solvent was removed *in vacuo* and the residue dissolved in ethyl acetate (10 mL). The organic phase was washed with brine (1 x 20 mL), dried over  $\text{MgSO}_4$ , filtered and the solvent removed *in vacuo*, affording a pale brown oil which was used without further purification (0.11 g, 100% crude).

**$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):**  $\delta$ /ppm 8.13 (1H, brs), 7.96 (2H, d,  $J = 8.7$ ), 7.82 (1H, brs), 7.62 (2H, d,  $J = 8.8$ ), 7.43-7.32 (5H, m), 6.37 (1H, d,  $J = 9.2$  Hz), 5.45 (1H, d,  $J = 9.3$  Hz), 5.08-4.90 (1H, m), 4.90-4.74 (1H, m), 3.92-3.58 (2H, m), 2.66-2.52 (1H, m), 2.20 (3H, s), 1.62-1.30 (2H, m), 1.30-1.05 (4H, m), 0.88 (9H, s), 0.78 (3H, t,  $J = 6.9$  Hz).  
**LRMS:** ( $m/z$ ) 510.4 ( $\text{M}+\text{H}$ ) $^+$ , 532.4 ( $\text{M}+\text{Na}$ ) $^+$ .

**STEP C:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-acetylamino benzoyl)-2',2'-dimethylpropyl)amide.

To a degassed solution of 2(*R*)-[(*N*-benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-(4-acetylamino benzoyl)-2',2'-dimethylpropyl)-amide (0.11 g, 0.21 mmol) in methanol (2 mL) was added 10% palladium on charcoal (20 mg). Hydrogen gas was bubbled through the suspension for 10 minutes when a balloon containing hydrogen was introduced to the reaction. After 2 hours, the catalyst was removed by filtration and the solvents were removed *in vacuo* to give a pale brown solid. Purification by reverse phase preparative HPLC afforded the title compound as a white crystalline solid (54 mg, 61%).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.30 (0.6 H, d, J = 9.1 Hz), 8.27 (0.4H, s), 8.19 (0.4H, d, J = 9.1 Hz), 7.99 (2H, d, J = 8.8 Hz), 7.86 (0.6H, s), 7.71 (2H, d, J = 8.8 Hz), 5.47 (1H, d, J = 9.1 Hz), 3.88-3.72 (1H, m), 3.61 (0.3H, dd, J = 13.9, 5.7 Hz), 3.41 (0.7H, dd, J = 14.1, 4.6 Hz), 3.09-2.95 (0.7H, m), 2.95-2.80 (0.3H, m), 2.15 (3H, s), 1.59-1.30 (2H, m), 1.30-1.04 (4H, m), 0.96 (9H, s), 0.74 (3H, t, J = 7.0 Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 200.4, 176.6, 176.3, 172.4, 159.8, 145.4, 134.8, 131.3, 120.6, 60.8, 53.9, 50.3, 45.37, 45.5, 36.2, 31.6, 30.6, 27.9, 24.5, 34.0, 14.5.


LRMS: (*m/z*) 420.2 (M+H)<sup>+</sup>, 442.2 (M+Na)<sup>+</sup>, 418.4 (M-H)<sup>-</sup>.

MIC (*E. coli*): 12.5 μM

MIC (*S. capitis*): 3.1 μM

### Example 23

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-methanesulfonylamino benzoyl)-2',2'-dimethylpropyl)amide



The title compound was prepared in an analogous manner to 2(*R*)-[[(*N*-formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-acetylamino)benzoyl)-2',2'-dimethylpropyl)amide as described in **Scheme 13**, using methanesulfonyl chloride in **Step B**. The relevant spectroscopic data are described below.

**STEP B:** 2(*R*)-[[(*N*-Benzyloxyl-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-(4-methanesulfonylamino)benzoyl)-2',2'-dimethylpropyl)-amide.

The title compound was afforded as a pale brown oil and used without further purification (0.10 g, 91% crude).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 8.29 (1H, s), 8.15 (0.5H, brs), 8.03-7.94 (2.5H, m), 7.82 (1H, brs), 7.62 (2H, d, *J* = 8.8), 7.45-7.32 (7H, m), 6.45 (1H, d, *J* = 9.2 Hz), 5.46 (1H, d, *J* = 9.3 Hz), 5.10-4.90 (1H, m), 4.90-4.76 (1H, m), 4.04-3.84 (1H, m), 3.82-3.62 (1H, m), 3.69 (3H, s), 2.70-2.57 (1H, m), 1.62-1.32 (2H, m), 1.32-1.05 (4H, m), 0.90 (9H, s), 0.77 (3H, t, *J* = 6.9 Hz).

**LRMS:** (*m/z*) 546.2 (*M*+H)<sup>+</sup>, 568.2 (*M*+Na)<sup>+</sup>.

**STEP C:** 2(*R*)-[[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-methanesulfonylamino)benzoyl)-2',2'-dimethylpropyl)amide

Purification by reverse phase preparative HPLC afforded the title compound as a pale brown crystalline solid (30 mg, 31%).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.30 (0.4 H, d, J = 9.2 Hz), 8.27 (0.4H, s), 8.19 (0.6H, d, J = 8.7 Hz), 8.02 (2H, d, J = 8.8 Hz), 7.86 (0.6H, s), 7.33 (2H, d, J = 8.8 Hz), 5.46 (1H, d, J = 9.0 Hz), 3.87-3.72 (1H, m), 3.61 (0.3H, dd, J = 14.1, 5.7 Hz), 3.42 (0.7H, dd, J = 14.2, 4.6 Hz), 3.06 (3H, s), 3.09-2.95 (0.7H, m), 2.95-2.80 (0.3H, m), 1.59-1.30 (2H, m), 1.30-1.02 (4H, m), 0.98 (3H, s), 0.97 (6H, s), 0.76 (3H, t, J = 7.0 Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 200.4, 200.3, 176.7, 176.4, 159.8, 145.2, 134.7, 131.9, 119.4, 60.8, 53.9, 50.3, 45.7, 45.4, 40.4, 36.2, 31.6, 30.6, 27.9, 24.0, 14.6.


LRMS: (m/z) 456.2 (M+H)<sup>+</sup>, 478.2 (M+Na)<sup>+</sup>, 454.4 (M-H)<sup>-</sup>.

MIC (*E. coli*): 50 μM


MIC (*S. capitis*): 3.1 μM

#### Example 24

2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)- (4-morpholin-4-ylbenzoyl)-propyl)amide



The title compound was prepared in as described in **Scheme 14**, followed by the **General Protocol (GP)**. The experimental details of the synthesis and relevant spectroscopic data are described below.



**Reagents and Conditions:**

A. Morpholine, DMSO, Et<sub>3</sub>N, 120°C

**Scheme 14**

**STEP A:** [2,2-Dimethyl-1(S)-(4-morpholin-4-ylbenzoyl)-propyl]-carbamic acid *tert*-butyl ester.

Morpholine (1.41 mL, 16.2 mmol) and triethylamine (0.25 mL, 1.8 mmol) were added to a solution of [1(S)-(4-fluorobenzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester (0.5 g, 1.62 mmol) in DMSO (8 mL), and the mixture heated at 120 °C for 15 hours. The solvent was removed *in vacuo* and the residual solid taken up in ethyl acetate (30 mL), and washed with 10% citric acid (aq.) (1 x 40 mL). The aqueous layer re-extracted with ethyl acetate (1 x 30 mL) and the combined organic phases were washed with brine (1 x 40 mL), dried over MgSO<sub>4</sub>, filtered and the solvent removed *in vacuo* to give a pale brown oil which was used without further purification (0.63 g, 100% crude).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.95 (2H, d, J = 9.0 Hz), 6.88 (2H, d, J = 9.0 Hz), 5.46 (1H, brd, J = 9.6 Hz), 5.10 (1H, d, J = 9.7 Hz), 3.90-3.82 (4H, m), 3.38-3.28 (4H, m), 1.42 (9H, s), 0.93 (9H, s).

LRMS: (m/z) 377.2 (M+H)<sup>+</sup>, 399.2 (M+Na)<sup>+</sup>.

**STEP GPA:** 2(S)-Amino-3,3-dimethyl-1-(4-morpholin-4-ylphenyl)-butan-1-one.

The title compound was recovered as pale brown solid and used without further purification (0.52 g, 100% crude).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.06 (2H, d, J = 9.0 Hz), 7.31 (2H, d, J = 9.0 Hz), 4.96 (1H, s), 3.99-3.89 (4H, m), 3.55-3.45 (4H, m), 1.04 (9H, s).

LRMS: (m/z) 277.2 (M+H)<sup>+</sup>, 299.2 (M+Na)<sup>+</sup>.

**STEP GPB:** 2(R)-[(N-Benzyloxy-N-formylamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)-(4-morpholin-4-ylbenzoyl)-propyl)-amide

Purification by column chromatography (SiO<sub>2</sub>, 2:1 to 2:3 *iso*-hexane:ethyl acetate) afforded the title compound as a pale brown oil (0.63 g, 72%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.13 (0.6H, brs), 8.98-7.88 (0.4, m), 7.94 (2H, d, J = 8.8 Hz), 7.48-7.30 (5H, m), 6.86 (2H, d, J = 9.0), 6.36 (1H, d, J = 9.3 Hz), 5.46 (1H, d, J = 9.4 Hz), 5.10-4.90 (1H, m), 4.90-4.75 (1H, m), 3.90-3.80 (4H, m), 3.90-3.53 (2H, m), 3.38-3.28 (4H, m), 2.62-2.50 (1H, m), 1.60-1.30 (2H, m), 1.30-1.02 (4H, m), 0.90 (9H, s), 0.76 (3H, t, J = 6.9 Hz).

LRMS: (m/z) 538.4 (M+H)<sup>+</sup>, 560.4 (M+Na)<sup>+</sup>, 536.4 (M-H)<sup>-</sup>.



**STEP GPC:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-(4-morpholin-4-ylbenzoyl)-propyl)amide

Purification by reverse phase preparative HPLC yielded the title compound as a pale beige foam (0.36 g, 69%).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.27 (0.3 H, s), 7.93 (2H, d, *J* = 9.0 Hz), 7.85 (0.7H, s), 6.97 (2H, d, *J* = 9.1 Hz), 5.46 (1H, s), 3.88-3.73 (5H, m), 3.61 (0.3H, dd, *J* = 13.8, 5.8 Hz), 3.41 (0.7H, dd, *J* = 14.2, 4.6 Hz), 3.39-3.27 (4H, m), 3.09-2.92 (0.7H, m), 2.92-2.78 (0.3H, m), 1.57-1.28 (2H, m), 1.28-1.00 (4H, m), 0.97 (3H, s), 0.95 (6H, s), 0.74 (3H, t, *J* = 7.0 Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 199.6, 176.5, 176.2, 19.8, 156.7, 132.3, 129.4, 114.8, 68.1, 60.2, 53.9, 50.3, 48.9, 45.8, 45.6, 36.3, 31.6, 30.6, 27.9, 24.0, 14.6.


LRMS: (*m/z*) 448.2 (*M*+H)<sup>+</sup>, 470.2 (*M*+Na)<sup>+</sup>, 446.4 (*M*-H)<sup>-</sup>.

MIC (*E. coli*): 6.25 μM

MIC (*S. capitis*): 3.1 μM

### Example 25

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-[4-(4-benzylpiperazin-1-yl)-benzoyl]-2',2'-dimethylpropyl)amide



The title compound was prepared utilising 1-benzylpiperazine in **Step A** as described in **Scheme 14**, and using the **General Protocol (GP)** thereafter. The relevant spectroscopic data are described below.

**STEP A:** [1(*S*)-[4-(4-Benzylpiperazin-1-yl)-benzoyl]-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester.

1-Benzylloxycarbonyl piperazine (0.45 g, 2.03 mmol) Purification by column chromatography (SiO<sub>2</sub>, 5:1 *iso*-hexane:ethyl acetate) afforded the title compound as a yellow oil (0.68 g, 82%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.91 (2H, d, J = 8.9 Hz), 7.38-7.28 (5H, m), 6.84 (2H, d, J = 9.1 Hz), 5.46 (1H, brd, J = 9.4 Hz), 5.10 (1H, d, J = 9.6 Hz), 3.56 (2H, s), 3.41-3.33 (4H, m), 3.62-3.55 (4H, m), 1.42 (9H, s), 0.93 (9H, s).

LRMS: (*m/z*) 466.2 (M+H)<sup>+</sup>, 488.2 (M+Na)<sup>+</sup>.

**STEP GPA:** 2(*S*)-1-[4-(4-Benzylpiperazin-1-yl)-phenyl]-3,3-dimethyl-butan-1-one dihydrochloride

The title compound was recovered as an orange foam and used without further purification (0.12 g, 100% crude).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.00 (2H, d, J = 9.0 Hz), 7.66-7.58 (2H, m), 7.54-7.47 (3H, m), 7.11 (2H, d, J = 9.0 Hz), 4.92 (1H, s), 4.44 (2H, s), 4.25-4.12 (2H, m), 3.62-3.51 (2H, m), 3.40-3.26 (4H, m), 1.03 (9H, s).

LRMS: (*m/z*) 366.2 (M+H)<sup>+</sup>, 388.2 (M+Na)<sup>+</sup>.

**STEP GPB:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-[4-(4-benzylpiperazin-1-yl)-benzoyl]-2',2'-dimethylpropyl)amide

Purification by column chromatography (SiO<sub>2</sub>, 2:1 ethyl acetate:iso-hexane) afforded the title compound as a pale yellow oil (37 mg, 24%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.13 (0.6H, brs), 7.98-7.87 (0.4, m), 7.93 (2H, d, J = 8.8 Hz), 7.48-7.30 (5H, m), 6.85 (2H, d, J = 9.0), 6.35 (1H, d, J = 9.3 Hz), 5.45 (1H, d, J = 9.3 Hz), 5.10-4.90 (1H, m), 4.90-4.75 (1H, m), 3.87-3.58 (2H, m), 3.56 (2H, s), 3.42-3.31 (4H, m), 2.63-2.50 (5H, m), 1.62-1.28 (2H, m), 1.28-1.05 (4H, m), 0.90 (9H, s), 0.76 (3H, t, J = 6.9 Hz).

LRMS: (*m/z*) 663.4 (M+Na)<sup>+</sup>.

**STEP GPC:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-[4-(4-benzylpiperazin-1-yl)-benzoyl]-2',2'-dimethylpropyl)amide trifluoroacetic acid

Purification by reverse phase preparative HPLC afforded the title compound as a colourless crystalline solid (10 mg, 27%).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.26 (0.4 H, s), 8.24 (0.3H, d, J = 9.8 Hz), 8.13 (0.3H, d, J = 9.1 Hz), 7.97 (2H, d, J = 8.9 Hz), 7.84 (0.6H, s), 7.59-7.48 (5H, m), 7.06 (2H, d, J = 8.9 Hz), 5.46 (0.5H, d, J = 5.6 Hz), 5.45 (0.5H, d, J = 9.1 Hz), 4.42 (2H, s), 3.79 (1H, dd, J=14.1, 8.8 Hz), 3.59 (1H, dd, J = 14.3, 5.8 Hz), 3.55-3.26 (8H, m), 3.10-2.95 (0.4H, m), 2.95-2.80 (0.6H, m), 1.56-1.31 (2H, m), 1.31-1.01 (4H, m), 0.96 (4H, s), 0.95 (5H, s), 0.75 (3H, t, J = 6.9).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 199.8, 176.3, 154.9, 132.8, 132.3, 131.9, 131.0, 130.9, 130.4, 116.0, 62.0, 60.5, 53.9, 52.7, 50.3, 46.3, 45.6, 45.5, 36.3, 31.6, 30.6, 27.9, 24.0, 14.6.


LRMS: (*m/z*) 533.4 (M+H)<sup>+</sup>, 559.4 (M+Na)<sup>+</sup>.

MIC (*E. coli*): 50  $\mu$ M

MIC (*S. capitis*): 3.1  $\mu$ M

### Example 26

2(*R*)-[*(N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)- (4-piperazin-1-yl-benzoyl)-propyl)amide



The title compound was prepared using 1-benzyloxycarbonylpiperazine in **Step A** as described in **Scheme 14**, and using the **General Protocol (GP)** thereafter. The experimental details of the synthesis and relevant spectroscopic data are described below.

**STEP A:** 4-[4-(2(*S*)-*tert*-Butoxycarbonylamino-3,3-dimethylbutyryl)-phenyl]-piperazine-1-carboxylic acid benzyl ester

Purification by column chromatography (SiO<sub>2</sub>, 5:1 *iso*-hexane:ethyl acetate) afforded the title compound as a yellow oil (0.68 g, 82%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 7.93 (2H, d, J = 8.9 Hz), 7.40-7.30 (5H, m), 6.85 (2H, d, J = 9.0 Hz), 5.42 (1H, brd, J = 8.9 Hz), 5.17 (2H, s), 5.10 (1H, d, J = 9.0 Hz), 3.70-3.62 (4H, m), 3.40-3.32 (4H, m), 1.42 (9H, s), 0.93 (9H, s).

**LRMS:** (*m/z*) 332.4 (M+Na)<sup>+</sup>.

**STEP GPA:** 4-[4-(2(*S*)-Amino-3,3-dimethylbutyryl)-phenyl]-piperazine-1-carboxylic acid benzyl ester hydrochloride

The title compound was recovered as a red foam and used without further purification (0.17 g, 100% crude).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):** δ/ppm 8.02 (2H, d, J = 9.0 Hz), 7.51 (2H, d, J = 9.0 Hz), 7.42-7.25 (5H, m), 5.17 (2H, s), 5.05 (1H, s), 3.89-3.79 (4H, m), 3.67-3.52 (4H, m), 1.03 (9H, s).

**LRMS:** (*m/z*) 410.2 (M+H)<sup>+</sup>, 432.2 (M+Na)<sup>+</sup>.

**STEP GPB:** 4-[4-(2(*S*)-{2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoylamino)-(3,3-dimethylbutyryl)-phenyl]-piperazine-1-carboxylic acid benzyl ester

Purification by column chromatography (SiO<sub>2</sub>, 3:1 *iso*-hexane:ethyl acetate) afforded the title compound as a pale yellow oil (0.15 g, 58%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 8.13 (0.6H, brs), 7.95-7.85 (0.4, m), 7.91 (2H, d, J = 8.9 Hz), 7.46-7.29 (5H, m), 6.84 (2H, d, J = 9.0), 6.38 (1H, d, J = 9.3 Hz), 5.45 (1H, d, J = 9.4 Hz), 5.17 (2H, s), 5.09-4.90 (1H, m), 4.90-4.75 (1H, m), 3.88-3.60 (6H, m), 3.42-3.31 (4H, m), 2.63-2.49 (1H, m), 1.62-1.28 (2H, m), 1.28-1.05 (4H, m), 0.90 (9H, s), 0.76 (3H, t, J = 7.0 Hz).

**LRMS:** (*m/z*) 671.2 (M+H)<sup>+</sup>, 693.4 (M+Na)<sup>+</sup>.

**STEP GPC:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-(4-piperazin-1-yl-benzoyl)-propyl)amide

The title compound was recovered as a yellow crystalline solid (0.12 g, 100%).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):** δ/ppm 8.27 (0.3H, s), 7.93 (2H, d, *J* = 9.0 Hz), 7.85 (0.7H, s), 6.99 (2H, d, *J* = 9.0 Hz), 5.46 (1H, s), 3.80 (1H, dd, *J* = 14.1, 9.7 Hz), 3.61 (0.5H, dd, *J* = 13.9, 5.7 Hz), 3.48-3.37 (4.5H, m), 3.08-2.96 (4.5H, m), 2.91-2.80 (0.5H, m), 1.57-1.32 (2H, m), 1.31-1.02 (4H, m), 0.95 (9H, s), 0.75 (3H, t, *J* = 7.0).

**<sup>13</sup>C NMR (CD<sub>3</sub>OD):** δ/ppm 199.6, 176.5, 176.2, 156.5, 132.3, 129.4, 115.1, 60.2, 54.0, 50.3, 48.36, 46.3, 45.8, 36.3, 31.6, 30.6, 27.0, 24.0, 14.6.


**LRMS:** (*m/z*) 447.2 (*M*+H)<sup>+</sup>, 469.2 (*M*+Na)<sup>+</sup>.

**MIC (*E. coli*):** 3.1 μM


**MIC (*S. capitis*):** 12.5 μM

### Example 27

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(4-methylpiperazin-1-yl)-benzoyl]-propyl)amide



The title compound was prepared from 4-[4-(2(*S*)-*tert*-butoxycarbonylamino-3,3-dimethylbutyryl)-phenyl]-piperazine-1-carboxylic acid benzyl ester as described in **Scheme 15**, and using the General Protocol (GP) thereafter. The experimental details of the synthesis and relevant spectroscopic data are described below.



**Reagents and Conditions:**

A. H<sub>2</sub>, 10% Pd-C, CH<sub>2</sub>O (37% wt. % in H<sub>2</sub>O), MeOH

**Scheme 15**

**STEP A:** {2,2-Dimethyl-1(*S*)-[4-(4-methylpiperazin-1-yl)-benzoyl]-propyl}-carbamic acid *tert*-butyl ester

Formaldehyde (37% wt. % solution in H<sub>2</sub>O, 0.5 mL) was added to a solution of 4-[4-(2(*S*)-*tert*-Butoxycarbonylamino-3,3-dimethylbutyryl)-phenyl]-piperazine-1-carboxylic acid benzyl ester in methanol (3 mL) and ethyl acetate (1 mL). The mixture was degassed and 10% palladium on charcoal (20 mg) was added. Hydrogen gas was bubbled through the suspension for 15 minutes when a balloon containing hydrogen was introduced to the reaction. After 18 hours, the catalyst was removed by filtration and the solvents were removed *in vacuo* to give a yellow oil which was used without further purification (0.15 g, 100% crude).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.97 (2H, d, J = 8.8 Hz), 7.40-7.30 (5H, m), 6.90 (2H, d, J = 8.9 Hz), 5.44 (1H, brd, J = 9.6 Hz), 5.09 (1H, d, J = 9.7 Hz), 3.87-3.78 (4H, m), 3.50-3.05 (4H, m), 2.85 (3H, s), 1.41 (9H, s), 0.92 (9H, s).

LRMS: (*m/z*) 390.2 (*M*+*H*)<sup>+</sup>, 402.4 (*M*+*Na*)<sup>+</sup>.

**STEP GPA:** 2(*S*)-Amino-3,3-dimethyl-1-[4-(4-methylpiperazin-1-yl)-phenyl]-butan-1-one.

The title compound was recovered as a yellow foam and used without further purification (0.14 g, 100% crude).

**STEP GPB:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(4-methylpiperazin-1-yl)-benzoyl]-propyl)amide trifluoroacetic acid. Purification by reverse phase preparative HPLC afforded the title compound as a yellow solid (0.15 g, 54%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.13 (0.6H, brs), 8.01-7.91 (0.4, m), 7.95 (2H, d, *J* = 8.7 Hz), 7.49-7.30 (5H, m), 6.90 (2H, d, *J* = 8.7), 6.40 (1H, d, *J* = 9.0 Hz), 5.44 (1H, d, *J* = 9.1 Hz), 5.10-4.90 (1H, m), 4.90-4.74 (1H, m), 4.00-3.33 (8H, m), 3.17-3.79 (2H, m), 2.89 (3H, s), 2.67-2.51 (1H, m), 1.62-1.30 (2H, m), 1.30-1.02 (4H, m), 0.90 (9H, s), 0.76 (3H, t, *J* = 7.0 Hz).

LRMS: (*m/z*) 551.2 (*M*+*H*)<sup>+</sup>, 573.4 (*M*+*Na*)<sup>+</sup>.

**STEP GPC:** 2(*R*)-[(*N*-Formyl-*N*-hydroxy-amino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(4-methylpiperazin-1-yl)-benzoyl]-propyl)amide trifluoroacetic acid.

Purification by reverse phase preparative HPLC afforded the title compound as a yellow solid (85 mg, 71%).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.27 (0.5H, s), 7.98 (2H, d, *J* = 8.9 Hz), 7.85 (0.5H, s), 7.08 (2H, d, *J* = 8.9 Hz), 5.46 (1H, s), 4.25-3.98 (2H, m), 3.80 (1H, dd, *J* = 14.2, 9.0 Hz), 3.71-3.51 (4H, m), 3.42 (1H, dd, *J* = 14.2, 4.5 Hz), 3.35-3.15 (4H, m), 3.11-2.92



(0.5H, m), 2.97 (3H, s), 2.92-2.81 (0.5H, m), 1.58-1.30 (2H, m), 1.30-1.03 (4H, m), 0.96 (3H, s), 0.95 (6H, s), 0.75 (3H, t, J = 9.9 Hz)

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 199.8, 176.6, 176.3, 154.9, 132.7, 131.0, 116.1, 60.5, 54.6, 53.9, 50.3, 46.5, 45.6, 45.5, 44.0, 36.3, 31.6, 30.3, 27.9, 24.0, 14.6.


LRMS: (m/z) 461.4 (M+H)<sup>+</sup>.

MIC (*E. coli*): 6.25 μM

MIC (*S. capitis*): 6.25 μM

### Example 28

2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)- [4-(4-pyrimidin-2-yl-piperazin-1-yl)-benzoyl]-propyl)amide



The title compound was prepared from 1-(2-pyrimidyl)-piperazine in **Step A** as described in **Scheme 14**, and using the **General Protocol (GP)** thereafter. The final step yielded both **Example 28** and **Example 29**. Relevant spectroscopic data are described below.

**STEP A:** {2,2-Dimethyl-1(*S*)-[4-(4-pyrimidin-2-ylpiperazin-1-yl)-benzoyl]-propyl}-carbamic acid *tert*-butyl ester.

Purification by column chromatography (SiO<sub>2</sub>, 5:2 *iso*-hexane:ethyl acetate) afforded the title compound as a pale brown oil (0.36 g, 82%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.34 (2H, d, J = 4.7), 7.95 (2H, d, J = 8.8 Hz), 6.90 (2H, d, J = 9.0 Hz), 6.54 (1H, t, J = 4.8), 5.48 (1H, brd, J = 9.4 Hz), 5.11 (1H, d, J = 9.7 Hz), 4.04-3.94 (4H, m), 3.54-3.43 (4H, m), 1.42 (9H, s), 0.94 (9H, s).

LRMS: (*m/z*) 454.4 (M+H)<sup>+</sup>.

**STEP GPA:** 2(*S*)-Amino-3,3-dimethyl-1-[4-(4-pyrimidin-2-ylpiperazin-1-yl)-phenyl]-butan-1-one dihydrochloride.

The title compound was recovered as pale brown solid and used without further purification (0.34 g, 100% crude).

<sup>1</sup>H NMR (CDOD<sub>3</sub>): δ/ppm 8.67 (2H, d, J = 5.3), 8.00 (2H, d, J = 9.1 Hz), 7.11-7.02 (3H, m), 4.90 (1H, s), 4.17-4.09 (4H, m), 3.82-3.72 (4H, m), 1.04 (9H, s).

LRMS: (*m/z*) 354.4 (M+H)<sup>+</sup>.

**STEP GPB:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(4-pyrimidin-2-ylpiperazin-1-yl)-benzoyl]-propyl)-amide

Purification by column chromatography (SiO<sub>2</sub>, 2:1 to 3:1 ethyl acetate:*iso*-hexane) afforded the title compound as a yellow foam (0.24 g, 83%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.13 (0.6H, brs), 7.98-7.88 (0.4, m), 7.94 (2H, d, J = 8.8 Hz), 7.48-7.30 (5H, m), 6.86 (2H, d, J = 9.0), 6.36 (1H, d, J = 9.3 Hz), 5.46 (1H, d, J = 9.4 Hz), 5.10-4.90 (1H, m), 4.90-4.75 (1H, m), 3.90-3.80 (4H, m), 3.90-3.53 (2H,

m), 3.38-3.28 (4H, m), 2.62-2.50 (1H, m), 1.60-1.30 (2H, m), 1.30-1.02 (4H, m), 0.90 (9H, s), 0.76 (3H, t, J = 6.9 Hz).

LRMS: (*m/z*) 538.4 (M+H)<sup>+</sup>, 560.4 (M+Na)<sup>+</sup>, 536.4 (M-H)<sup>-</sup>.

**STEP GPC:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(*S*)-[4-(4-pyrimidin-2-yl-piperazin-1-yl)-benzoyl]-propyl)amide

Purification by reverse phase preparative HPLC yielded the title compound as a pale yellow glassy solid (65 mg, 34%, 58% combined yield with **Example 29**).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.34 (2H, d, J = 4.8 Hz), 8.27 (0.3H, s), 7.95 (2H, d, J = 9.0 Hz), 7.86 (0.7H, s), 7.00 (2H, d, J = 9.0 Hz), 6.61 (1H, t, J = 4.8 Hz), 5.48 (1H, s), 3.95 (4H, t, J = 5.3 Hz), 3.87-3.73 (1H, m), 3.61 (0.4H, dd, J = 14.0, 5.8 Hz), 3.50 (4H, t, J = 5.3 Hz), 3.41 (0.6H, dd, J = 14.0, 4.6Hz), 3.09-2.95 (0.6H, m), 2.95-2.78 (0.4H, m), 1.58-1.30 (2H, m), 1.28-1.04 (4H, m), 0.96 (9H, s), 0.74 (3H, t, J = 7.0 Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 199.5, 176.5, 176.2, 163.2, 159.5, 156.4, 132.4, 129.0, 114.9, 111.9, 60.2, 56.9, 50.3, 45.8, 45.6, 44.9, 36.4, 31.6, 30.6, 28.0, 24.0, 14.6.


LRMS: (*m/z*) 525.2 (M+H)<sup>+</sup>, 547.2 (M+Na)<sup>+</sup>.

MIC (*E. coli*): 12.5 μM

MIC (*S. capitis*): 1.6 μM

**Example 29**

2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)- {4-[4-(1,4,5,6-tetrahydropyrimidin-2-yl)-piperazin-1-yl]-benzoyl}-propyl)amide



The title compound was prepared in an identical manner to **Example 28** as described in **Scheme 14**. The final step yielded both **Example 28** and **Example 29**. Relevant spectroscopic data are described below.

**STEP GPC:** 2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)- {4-[4-(1,4,5,6-tetrahydropyrimidin-2-yl)-piperazin-1-yl]-benzoyl}-propyl)amide.

Purification by reverse phase preparative HPLC yielded the title compound as a colourless glassy solid (47 mg, 24%, 58% combined yield with **Example 28**).

<sup>1</sup>H NMR (CD<sub>3</sub>OD): δ/ppm 8.27 (0.3 H, s), 7.95 (2H, d, J = 8.9 Hz), 7.86 (0.7H, s), 6.98 (2H, d, J = 8.9 Hz), 5.46 (1H, s), 3.77 (1H, dd, J = 14.1, 4.9 Hz), 3.69-3.50 (8.3H, m), 3.47-3.38 (0.7H, m), 3.42 (4H, t, J = 5.9 Hz), 3.10-2.95 (0.7H, m), 2.95-

2.80 (0.3H, m), 1.97 (2H, tt, J = 5.9, 5.9 Hz), 1.58-1.30 (2H, m), 1.30-1.02 (4H, m), 0.97 (3H, s), 0.95 (6H, s), 0.74 (3H, t, J = 7.0 Hz).

**<sup>13</sup>C NMR (CD<sub>3</sub>OD):** δ/ppm 199.8, 199.6, 176.5, 176.2, 159.8, 155.5, 155.3, 132.4, 129.5, 114.8, 60.3, 53.9, 50.3, 47.4, 46.1, 45.7, 45.6, 40.7, 36.3, 31.6, 30.6, 28.0, 24.0, 21.7, 14.6.


**LRMS:** ( $m/z$ ) 529.2 ( $M+H$ )<sup>+</sup>, 527.4 ( $M-H$ )<sup>-</sup>.

**MIC (*E. coli*): 50  $\mu$ M**

**MIC (*S. capitis*): 6.25  $\mu$ M**

### Example 30

2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(1'(S)-(4-dipropylaminobenzoyl)-2',2'-dimethylpropyl)amide



The title compound was prepared utilising diallyl amine in **Step A** as described in **Scheme 14**, and using the **General Protocol (GP)** thereafter. The final step yielded both **Example 30** and **Example 31**. Relevant spectroscopic data are described below.

**STEP A:** [1(*S*)-(4-Diallylamino benzoyl)-2,2-dimethylpropyl]-carbamic acid *tert*-butyl ester.

Purification by column chromatography (SiO<sub>2</sub>, 19:1 *iso*-hexane:ethyl acetate) afforded the title compound as a thick yellow oil (5.18 g, 52% (81% based on recovered SM)).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.89 (2H, d, J = 8.6 Hz), 6.66 (2H, d, J = 8.7 Hz), 5.85 (2H, ddt, J = 17.1, 10.3, 4.7 Hz), 5.47 (1H, brd, J = 9.5 Hz), 5.26-5.11 (4H, m), 5.09 (1H, d, J = 9.6 Hz), 4.04-3.94 (4H, m), 1.42 (9H, s), 0.95 (9H, s).

LRMS: (*m/z*) 387.2 (M+H)<sup>+</sup>.

**STEP GPA:** 2(*S*)-Amino-1-(4-diallylamino-phenyl)-3,3-dimethylbutan-1-one hydrochloride.

The title compound was recovered as yellow solid and used without further purification (3.7 g, 100% crude).

<sup>1</sup>H NMR (CDOD<sub>3</sub>): δ/ppm 7.91 (2H, d, J = 8.6 Hz), 6.85 (2H, d, J = 8.7 Hz), 5.90 (2H, ddt, J = 17.1, 10.3, 4.7 Hz), 5.26-5.17 (3H, m), 5.16-5.12 (1H, m), 4.84 (1H, s), 4.13-3.05 (4H, m), 1.05 (9H, s).

LRMS: (*m/z*) 287.2 (M+H)<sup>+</sup>.

**STEP GPB:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-(4-diallylamino benzoyl)-2',2'-dimethylpropyl)-amide

Purification by column chromatography (SiO<sub>2</sub>, 3:1 *iso*-hexane:ethyl acetate) afforded the title compound as a yellow oil (5.64 g, 90%).

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 8.13 (0.6H, brs), 7.95-7.83 (0.4, m), 7.88 (2H, d, J = 9.0 Hz), 7.48-7.31 (5H, m), 6.65 (2H, d, J = 9.1), 6.40 (1H, d, J = 9.3 Hz), 5.84 (2H, ddt, J = 17.0, 10.4, 4.7 Hz), 5.44 (1H, d, J = 9.4 Hz), 5.24-5.17 (3H, m), 5.16-5.10 (1H, m), 5.08-4.90 (1H, m), 4.90-4.74 (1H, m), 4.04-3.94 (4H, m), 3.89-3.58 (2H, m), 2.62-2.48 (1H, m), 1.60-1.30 (2H, m), 1.30-1.05 (4H, m), 0.91 (9H, s), 0.76 (3H, t, J = 6.9 Hz).

**LRMS:** (*m/z*) 548.4 (M+H)<sup>+</sup>, 570.4 (M+Na)<sup>+</sup>.

**STEP GPC:** 2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-(4-dipropylaminobenzoyl)-2',2'-dimethylpropyl)amide

Purification by reverse phase preparative HPLC yielded the title compound as a colourless foam (20 mg, 22%, 69% combined yield with **Example 31**).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):** δ/ppm 8.27 (0.3H, s), 7.92-7.80 (2.7H, m), 6.68 (2H, d, J = 9.1 Hz), 5.45 (1H, s), 3.88-3.75 (1H, m), 3.61 (0.3H, dd, J = 14.4, 5.8 Hz), 3.48-3.30 (4.7H, m), 3.08-2.90 (0.7H, m), 3.90-2.78 (0.3H, m), 1.64 (4H, qt, J = 7.5, 7.5 Hz), 1.55-1.27 (2H, m), 1.27-1.06 (4H, m), 1.05-0.76 (15H, m), 0.74 (3H, t, J = 6.9 Hz).

**<sup>13</sup>C NMR (CD<sub>3</sub>OD):** δ/ppm 198.8, 176.4, 176.1, 159.8, 154.2, 132.8, 125.8, 112.1, 59.9, 54.0, 50.3, 45.9, 45.7, 36.4, 31.6, 31.2, 30.6, 27.6, 24.0, 21.8, 14.5, 11.9


**LRMS:** (*m/z*) 462.2 (M+H)<sup>+</sup>, 460.4 (M-H)<sup>-</sup>.

**MIC (*E. coli*):** 100 μM

**MIC (*S. capitis*):** 3.1 μM

**Example 31**

2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)-(4-propylaminobenzoyl)-propyl)amide



The title compound was prepared in an identical manner to **Example 30** as described in **Scheme 14**. The final step yielded both **Example 30** and **Example 31**. Relevant spectroscopic data are described below.

**STEP GPC:** 2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(2',2'-dimethyl-1'(S)-(4-propylaminobenzoyl)-propyl)amide

Purification by reverse phase preparative HPLC yielded the title compound as a white powder (45 mg, 49%; 69% combined yield with **Example 30**).

**<sup>1</sup>H NMR (CD<sub>3</sub>OD):**  $\delta$ /ppm 8.27 (0.4 H, s), 7.89-7.78 (2.6H, m), 6.60 (2H, d, J=8.9Hz), 5.44 (1H, s), 3.86-3.72 (1H, m), 3.61 (0.4H, dd, J=14.0,5.8Hz), 3.40 (0.6H, dd, J=14.2,4.6Hz), 3.13 (2H, t, J=7.1Hz), 3.08-2.90 (0.6H, m), 3.90-2.76 (0.4H, m), 1.65 (2H, qt, J=7.3,7.3Hz), 1.55-1.30 (2H, m), 1.29-1.06 (4H, m), 1.04-0.89 (12H, m), 0.75 (3H, t, J = 7.0).



<sup>13</sup>C NMR (CD<sub>3</sub>OD): δ/ppm 199.1, 198.9, 176.4, 176.1, 159.8, 155.9, 132.9, 126.7, 112.5, 59.8, 53.9, 50.3, 46.1, 45.9, 45.7, 36.4, 31.6, 30.6, 28.0, 24.1, 24.0, 14.5, 12.3.


LRMS: (*m/z*) 420.2 (M+H)<sup>+</sup>, 442.2 (M+Na)<sup>+</sup>, 418.4 (M-H)<sup>-</sup>.

MIC (*E. coli*): 12.5 μM

MIC (*S. capitis*): 3.1 μM

### Example 32

2(*R*)-[(*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-acetyl-2',2'-dimethylpropyl)amide



#### STEP A: 2(*S*)-*tert*-Butoxycarbonylamino-3,3-dimethyl-butyric acid

To a suspension of L-*tert*-leucine (26.1 g, 0.2 mol) in methanol (150 mL) was added triethylamine (56 mL, 0.4 mol) and the mixture cooled to 0 °C. To the mixture was added slowly a solution of di-*tert*-butyldicarbamate (48 g, 0.22 mol) in methanol (40 mL) such that an internal temperature of between 0 and 5 °C was maintained. The reaction was allowed to stir overnight and the solvents were removed *in vacuo*. The residue was dissolved in ethyl acetate (200 mL) and washed with 10% w/v aqueous citric acid solution (3 x 100 mL). The organic layer was dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give the title product as a pale yellow oil (48.9 g, >100 %, residual solvent).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 9.20 (1H, bs), 5.10 (1H, m), 4.15 (1H, m), 1.45 (9H, s), 1.00 (9H, s).

**STEP B:** [1(S)-(Methoxy-methyl-carbamoyl)-2,2-dimethylpropyl]-carbamic acid-*tert*-butyl ester.

To a solution of 2(S)-*tert*-butoxycarbonylamino-3,3-dimethyl-butyric acid (30 g, 0.13 mol) in dichloromethane (400 mL) was added portionwise 1,1'-carbonyldiimidazole (63.3 g, 0.39 mol). After addition was complete the reaction was allowed to stir for 15 minutes before *N*-methoxy-*N*-methylamine hydrochloride (38 g, 0.39 mol) was added and the reaction allowed to stir overnight. The solvents were removed *in vacuo* and the residue partitioned between ethyl acetate (250 mL) and 1M HCl (150 mL). The organic phase was washed with 1M HCl solution (2 x 100 mL), saturated sodium bicarbonate solution (2 x 100 mL) and brine (100 mL). The organic phase was dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 30% ethyl acetate in hexanes) afforded the title product as a pale yellow oil (26.5 g, 74 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 5.20 (1H, bd, J = 9.5 Hz), 4.65 (1H, d, J = 10.4 Hz), 3.77 (3H, s), 3.21 (3H, s), 1.43 (9H, s), 0.97 (9H, s).

LRMS: (*m/z*) 297 (M+Na)<sup>+</sup>.

**STEP C:** (1(S)-Acetyl-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester

[1(S)-(Methoxy-methyl-carbamoyl)-2,2-dimethylpropyl]-carbamic acid-*tert*-butyl ester (2.0 g, 7.3 mmol) was dissolved in freshly distilled THF under an argon atmosphere. The solution was cooled to -78 °C whereupon MeLi (15.6 mL, 1.4 M in diethyl ether, 21.9 mmol, 3 eqv) was added dropwise maintaining an internal temperature < -70 °C. The reaction was stirred at -70 °C for 30 minutes before warming to -30 °C for a

further 30 minutes. After this time TLC analysis indicated the complete consumption of starting material. Saturated aqueous ammonium chloride solution (10 mL) was added and the reaction allowed to warm to room temperature. The aqueous phase was extracted with ethyl acetate (3 x 30 mL). The organic phases were washed with water (3 x 50 mL) and brine (50 mL), dried over  $\text{MgSO}_4$ , filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography ( $\text{SiO}_2$ , 20 % ethyl acetate in hexanes) afforded the title compound as a colourless oil (1.355 g, 84%).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$ /ppm 5.13 (1H, m), 4.16 (1H, bd,  $J = 10.0$  Hz), 2.25 (3H, s), 1.43 (9H, s), 0.98 (9H, s).

LRMS: ( $m/z$ ) 252 ( $\text{M}+\text{Na}$ ) $^+$ .

#### STEP D: 3(S)-Amino-4,4-dimethyl-pentan-2-one

To a solution of (1(S)-acetyl-2,2-dimethylpropyl)carbamic acid-*tert*-butyl ester (700 mg, 2.55 mmol) in dichloromethane (10 mL) at 0 °C was added trifluoroacetic acid (5 mL). The reaction was left at <5 °C for 16 hours before the solvents were removed *in vacuo* to give a bright yellow oil. The residue was dissolved in methanol (30 mL) and was treated portionwise with Dowex 1X8-400 basic resin until pH 9. Filtration and removal of the solvents *in vacuo* gave the title compound as a pale yellow oil (~1.0 g, > 100 %, residual water). The material was used without further purification.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$ /ppm 3.28 (1H, s), 2.21 (3H, s), 1.50 (2H, bs), 0.98 (9H, s).

#### STEP E: 2(R)-(N-Benzyloxyaminomethyl)-hexanoic acid-(1'(S)-acetyl-2',2'-dimethylpropyl)amide

To a suspension of 2(R)-(N-benzyloxyamino-methyl)-hexanoic acid (879 mg, 3.5 mmol, 1.1 eqv) in DMF (10 mL) at 0 °C was added EDC (730 mg, 1.2 eqv) and 1-

hydroxybenzotriazole (43 mg, 10 mol%). The mixture was allowed to stir at 0 °C for 45 minutes before a solution of 3(*S*)-amino-4,4-dimethyl-pentan-2-one (409 mg, 1 eqv) in DMF (5 mL) was added. The reaction was allowed to warm to room temperature and stirred overnight. The solvents were removed *in vacuo* and the residue dissolved in ethyl acetate (50 mL) and 10% w/v aqueous citric acid (50 mL). The aqueous phase was extracted with ethyl acetate (3 x 50 mL) and the combined organic layers washed with saturated sodium hydrogen carbonate (100 mL) and brine (2 x 100 mL). The organic phases were dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 25% ethyl acetate in hexanes) gave the title compound as a clear oil (412 mg, 36 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.35 (5H, m), 6.73 (1H, bd, J = 8.7 Hz), 5.68 (1H, bs), 4.73 (2H, s), 4.54 (1H, d, J = 8.7 Hz), 3.06 (2H, m), 2.50 (1H, m), 2.27 (3H, s), 1.70 - 1.20 (6H, m), 0.98 (9H, s), 0.87 (3H, t, J = 7.0 Hz).

LRMS: (*m/z*) 363 (M+H)<sup>+</sup>.

**STEP F:** 2(*R*)-[*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-acetyl-2',2'-dimethylpropyl)amide

To a solution of 2(*R*)-(*N*-benzyloxyamino-methyl)-hexanoic acid-(1'(*S*)-acetyl-2',2'-dimethylpropyl)amide (400 mg, 1.1 mmol) in dichloromethane (10 mL) at 0 °C was added formylacetic anhydride (0.25 mL, 2.76 mmol, 2.5 eqv) and the reaction allowed to stir at room temperature. After 2 hours TLC analysis indicated complete consumption of the starting material. The solvents were removed *in vacuo* and the residue azeotroped with toluene (3 x 30 mL). The product was placed under high vacuum to give the title compound as a clear sticky oil (443 mg, >100 % , residual solvent).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 8.15 (0.7H, bs), 7.90 (0.3H, bs), 7.38 (5H, bs), 6.06 (1H, bd,

$J = 8.2$  Hz), 5.00 - 4.75 (2H, m), 4.43 (1H, bd,  $J = 8.3$  Hz), 3.72 (2H, m), 2.53 (1H, m), 2.24 (3H, s), 1.65 - 1.40 (2H, m), 1.35 - 1.10 (4H, m), 0.94 (9H, s), 0.86 (3H, t,  $J = 6.7$  Hz).

**LRMS:** ( $m/z$ ) 389 (M-H)<sup>-</sup>, 391 (M+H)<sup>+</sup>.

**STEP G:** 2(*R*)-[*N*-Formyl-*N*-hydroxyamino)-methyl]-hexanoic acid-(1'(*S*)-acetyl-2',2'-dimethylpropyl)amide

To a solution of 2(*R*)-[*N*-benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-acetyl-2',2'-dimethylpropyl)amide (200 mg, 0.55 mmol) in methanol (10 mL) under argon was added a slurry of 10% Pd/C (27 mg) in ethyl acetate (5 mL). Hydrogen gas was bubbled through the reaction for 15 minutes after which time the reaction was allowed to stir under a hydrogen atmosphere of 1 bar for two hours. After this time TLC analysis indicated the complete consumption of starting material. The reaction was flushed with argon and filtered through Celite. The solvents were removed *in vacuo* to afford the title compound as a pale orange solid (148 mg, 96%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$ /ppm 8.38 (0.3H, s), 7.81 (0.7H, s), 6.40 (0.3H, bd,  $J = 6.9$  Hz), 6.25 (0.7H, bd,  $J = 8.2$  Hz), 4.52 (1H, m), 4.00 (0.3H, dd,  $J = 14.7, 7.1$  Hz), 3.82 (0.7H, dd,  $J = 14.2, 10.0$  Hz), 3.62 (0.3H, dd,  $J = 14.5, 2.8$  Hz), 3.47 (0.7H, dd,  $J = 14.2, 3.8$  Hz), 2.79 (0.7H, m), 2.65 (0.3H, m), 2.30 (0.9H, s), 2.27 (2.1H, s), 1.60 (1H, m), 1.50 - 1.20 (5H, m), 1.01 (2.7H, s), 0.96 (6.3H, s), 0.88 (3H, t,  $J = 6.6$  Hz).

<sup>13</sup>C NMR (CD<sub>3</sub>OD):  $\delta$ /ppm 177.0, 176.7, 67.9, 53.9, 45.5, 45.3, 35.1, 31.5, 31.2, 30.7, 27.8, 24.1, 14.6.

**LRMS:** ( $m/z$ ) 299 (M-H)<sup>-</sup>, 301 (M+H)<sup>+</sup>, 323 (M+Na)<sup>+</sup>.

MISSING AT THE TIME OF PUBLICATION

To a suspension of 2(*R*)-(N-benzyloxyamino-methyl)-hexanoic acid (1.430 mg, 5.17 mmol, 1.1 eqv) in DMF (20 mL) at 0 °C was added EDC (1.190 g, 5.64 mmol 1.2 eqv) and 1-hydroxybenzotriazole (84 mg, 10 mol%). The mixture was allowed to stir at 0 °C for 45 minutes before a solution of 2(*S*)-amino-N-methoxy-3,3,*N*-trimethylbutyramide (900 mg, 4.7 mmol, 1 eqv) in DMF (5 mL) was added. The reaction was allowed to warm to room temperature and stirred overnight. The solvents were removed *in vacuo* and the residue dissolved in ethyl acetate (75 mL) and 10% w/v aqueous citric acid (50 mL). The aqueous phase was extracted with ethyl acetate (3 x 75 mL) and the combined organic layers washed with saturated sodium hydrogen carbonate solution (100 mL) and brine (2 x 100 mL). The organic phases were dried over MgSO<sub>4</sub>, filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography (SiO<sub>2</sub>, 25% ethyl acetate in hexanes) gave the title compound as a clear oil (1.43 g, 68 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub>): δ/ppm 7.35 (5H, m), 6.73 (1H, bd, J = 10.0 Hz), 5.70 (1H, bs), 5.05 (1H, d, J = 10.0 Hz), 4.74 (2H, s), 3.78 (3H, s), 3.20 (3H, s), 3.07 (2H, m), 2.50 (1H, m), 1.65 (1H, m), 1.45 (1H, m), 1.25 (4H, m), 0.98 (9H, s), 0.86 (3H, t, J = 7.0 Hz).

LRMS: (*m/z*) 408 (M+H)<sup>+</sup>.

**STEP C:** 2(*R*)-[(N-Benzyloxyamino)-methyl]-hexanoic acid-(1'(*S*)-benzoyl-2',2'-dimethylpropyl)amide

To a solution of 2(*R*)-[(N-benzyloxyamino)-methyl]-hexanoic acid-[1'(*S*)-(methoxymethyl-carbamoyl)-2',2'-dimethylpropyl]amide (416 mg, 1.05 mmol) in freshly distilled THF (10 mL) under argon at -78 °C was added dropwise PhLi (3.5 mL, 1.8M in cyclohexane, 6.3 mmol, 6 eqv) such that an internal temperature of < -70 °C was maintained. The reaction was allowed to stir at -70 °C for three hours after which time TLC analysis appeared to indicate the complete consumption of starting material. Saturated aqueous ammonium chloride solution (10 mL) was added and

the solution allowed to warm to room temperature. The aqueous phase was extracted with ethyl acetate (3 x 50 mL), and the organic phases combined and washed with water (3 x 50 mL) and brine (50 mL). The organic phase was dried over  $\text{MgSO}_4$ , filtered and the solvents removed *in vacuo* to give a crude yellow oil. Purification by column chromatography ( $\text{SiO}_2$  / 25 % ethyl acetate in hexanes) gave the title product as a colourless oil (143 mg, 32 %,  $R_f = 0.3$ ) and the recovered starting material 2(*R*)-[(*N*-Benzyloxyamino)-methyl]-hexanoic acid-[1'(*S*)-(methoxymethyl-carbamoyl)-2',2'-dimethylpropyl]amide (209 mg,  $R_f = 0.1$ ).

$^1\text{H NMR}$  ( $\text{CDCl}_3$ ):  $\delta$ /ppm 8.02 (2H, m), 7.62 - 7.40 (2H, m), 7.40 - 7.25 (6H, m), 6.97 (1H, bd,  $J = 9.4$  Hz), 5.60 (1H, d,  $J = 9.4$  Hz), 4.75 (2H, s), 3.10 (2H, m), 2.53 (1H, m), 1.65 - 1.30 (2H, m), 1.30 - 1.10 (4H, m), 0.94 (9H, s), 0.85 (3H, m).

**STEP D:** 2(*R*)-[(*N*-Benzyloxy-*N*-formylamino)-methyl]-hexanoic acid-(1'(*S*)-benzoyl-2',2'-dimethylpropyl)amide

To a solution of 2(*R*)-[(*N*-benzyloxyamino)-methyl]-hexanoic acid-(1'(*S*)-benzoyl-2',2'-dimethylpropyl)amide (195 mg, 0.46 mmol) in dichloromethane (50 mL) at 0 °C was added formylacetic anhydride (0.1 mL, 1.15 mmol, 2.5 eqv) and the reaction allowed to stir at room temperature. After 1 hour TLC analysis indicated complete consumption of the starting material. The solvents were removed *in vacuo* and the residue azeotroped with toluene (3 x 30 mL). The product was placed under high vacuum for several hours to give the title compound as a clear sticky oil (208 mg, 100 %).

$^1\text{H NMR}$  ( $\text{CDCl}_3$ ):  $\delta$ /ppm 8.15 (0.5H, bs), 8.00 (2.5H, m), 7.65 - 7.30 (8H, m), 6.33 (1H, bd,  $J = 9.0$  Hz), 5.51 (1H, d,  $J = 9.0$  Hz), 5.00 (1H, m), 4.80 (1H, m), 3.75 (2H, bm), 2.58 (1H, m), 1.70 - 1.00 (6H, m), 0.89 (9H, s), 0.77 (3H, t,  $J = 6.7$  Hz).

LRMS: ( $m/z$ ) 451 ( $\text{M-H}$ ) $^+$ , 453 ( $\text{M+H}$ ) $^+$ .



**STEP E: 2(R)-[(N-Formyl-N-hydroxyamino)-methyl]-hexanoic acid-(1'(S)-benzoyl-2',2'-dimethylpropyl)amide**

To a solution of 2(R)-[(N-benzyloxy-N-formylamino)-methyl]-hexanoic acid-(1'(S)-benzoyl-2',2'-dimethylpropyl)amide (192 mg, 0.43 mmol) in methanol (5 mL) was added a slurry of 10% Pd/C (20 mg) in ethyl acetate (2 mL). Hydrogen gas was bubbled through the reaction for 15 minutes after which time the reaction was allowed to stir under a hydrogen atmosphere of 1 bar for. After four hours reaction time TLC analysis indicated the complete consumption of starting material. The reaction was flushed with argon and filtered through glass fibre. The solvents were removed *in vacuo* to afford the title compound as a white crystalline solid (135 mg, 88%). The product was purified by preparative HPLC giving 75 mg (49 %) of a white crystalline solid.

**<sup>1</sup>H NMR (CDCl<sub>3</sub>):** δ/ppm 9.18 (1H, bs), 8.40 (0.3H, s), 8.00 (2H, bd, J = 7.1 Hz), 7.86 (0.7H, s), 7.60 (1H, m), 7.48 (2H, m), 6.84 (0.3H, bd, J = 9.2 Hz), 6.70 (0.7H, bd, J = 9.3 Hz), 5.55 (1H, m), 4.05 (0.3H, dd, J = 14.6, 7.2 Hz), 3.87 (0.7H, dd, J = 14.1, 9.7 Hz), 3.64 (0.3H, dd, J = 14.6, 3.3 Hz), 3.47 (0.7H, dd, J = 14.0, 4.0 Hz), 2.85 (0.7H, m), 2.71 (0.3H, m), 1.55 (1H, m), 1.40 (1H, m), 1.30 - 1.10 (4H, m), 0.94 (3H, s), 0.90 (6H, s), 0.78 (3H, t, J = 6.3 Hz).

**<sup>13</sup>C NMR (CDCl<sub>3</sub>):** δ/ppm 176.1, 172.9, 137.6, 133.9, 133.6, 128.9, 128.8, 128.6, 59.4, 59.1, 51.8, 47.7, 46.2, 44.7, 35.7, 35.4, 30.0, 29.8, 29.2, 29.1, 27.0, 22.5, 13.7.


**LRMS:** (*m/z*) 361 (M-H)<sup>-</sup>, 363 (M+H)<sup>+</sup>.

Biological Example

Minimal inhibitory concentrations (MIC) of inhibitors against *E. coli* strain DH5 $\alpha$  (Genotype; F- $\phi$ 80d/*lacZ* $\Delta$ M15 $\Delta$ (*lacZYA-argF*)U169 *deoR recA1 endA1 hsdR17*( $r_k^-$ ,  $m_k^+$ )/*phoA supE44* $\lambda^-$  *thi-1 gyrA96 relA1*) obtained from GibcoBRL Life Technologies, or *Staphylococcus capitis* (American Type Culture Collection number 35661) were determined as follows. Stock solutions of each test compound were prepared by dissolution of the compound in dimethylsulfoxide at 10mM. For the determination of the minimal inhibitory concentration, two fold serial dilutions were prepared in 2xYT broth (typtone 16g/1, yeast extract 10g/1, sodium chloride 5g/1 obtained from BIO 101 Inc, 1070 Joshua Way, Vista, CA92083, USA) to yield 0.05 ml compound-containing medium per well. Inocula were prepared from cultures grown overnight in 2xYT broth at 37°C. Cell densities were adjusted to absorbance at 660nm ( $A_{660}$ ) = 0.1; the optical density-standardised preparations were diluted 1:1000 in 2xYT broth; and each well inoculated with 0.05ml of the diluted bacteria. Microtiter plates were incubated at 37°C for 18 hours in a humidified incubator. The MIC ( $\mu$ M) was recorded as the lowest drug concentration that inhibited visible growth.

## Claims

1. The use of a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt, hydrate or solvate thereof in the preparation of a composition for treatment of bacterial or protozoal infections in humans and non-human mammals:



wherein:

Z represents a radical of formula  $-\text{N}(\text{OH})\text{CH}(=\text{O})$  or formula  $-\text{C}(=\text{O})\text{NH}(\text{OH})$ ;

$\text{R}_1$  represents hydrogen, methyl or trifluoromethyl, or, except when Z is a radical of formula  $-\text{N}(\text{OH})\text{CH}(=\text{O})$ , a hydroxy or amino group;

$\text{R}_2$  represents a group  $\text{R}_{10}-(\text{X})_n-(\text{ALK})_m$ - wherein

$\text{R}_{10}$  represents hydrogen, or a  $\text{C}_1$ - $\text{C}_6$  alkyl,  $\text{C}_2$ - $\text{C}_6$  alkenyl,  $\text{C}_2$ - $\text{C}_6$  alkynyl, cycloalkyl, aryl, or heterocyclyl group, any of which may be unsubstituted or substituted by  $(\text{C}_1$ - $\text{C}_6)$ alkyl,  $(\text{C}_1$ - $\text{C}_6)$ alkoxy, hydroxy, mercapto,  $(\text{C}_1$ - $\text{C}_6)$ alkylthio, amino, halo (including fluoro, chloro, bromo and iodo), trifluoromethyl, cyano, nitro, oxo,  $-\text{COOH}$ ,  $-\text{CONH}_2$ ,  $-\text{COOR}^A$ ,  $-\text{NHCOR}^A$ ,  $-\text{CONHR}^A$ ,  $-\text{NHR}^A$ ,  $-\text{NR}^A\text{R}^B$ , or  $-\text{CONR}^A\text{R}^B$  wherein  $\text{R}^A$  and  $\text{R}^B$  are independently a  $(\text{C}_1$ - $\text{C}_6)$ alkyl group and

ALK represents a straight or branched divalent  $\text{C}_1$ - $\text{C}_6$  alkylene,  $\text{C}_2$ - $\text{C}_6$  alkenylene, or  $\text{C}_2$ - $\text{C}_6$  alkynylene radical, and may be interrupted by one or more non-adjacent  $-\text{NH}-$ ,  $-\text{O}-$  or  $-\text{S}-$  linkages,

X represents  $-\text{NH}-$ ,  $-\text{O}-$  or  $-\text{S}-$ , and

m and n are independently 0 or 1;

R<sub>3</sub> represents the side chain of a natural or non-natural alpha amino acid; and

R<sub>4</sub> represents a radical R<sub>5</sub>-(ALK)<sub>p</sub>, wherein ALK is as defined in relation to R<sub>2</sub>, p is 0 or 1, and R<sub>5</sub> represents hydrogen, or a C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, cycloalkyl, aryl, or heterocyclyl group any of which

(i) may be substituted by a group selected from (C<sub>1</sub>-C<sub>6</sub>)alkyl, phenyl, benzyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, phenoxy, hydroxy, mercapto, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, amino, halo, trifluoromethyl, cyano, nitro, oxo, -COOH, -SO<sub>2</sub>H, -CONH<sub>2</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -COR<sup>A</sup>, -SOR<sup>A</sup>, -SO<sub>2</sub>R<sup>A</sup>, -SO<sub>2</sub>R<sup>A</sup>, -COOR<sup>A</sup>, -CONHR<sup>A</sup>, -SO<sub>2</sub>NHR<sup>A</sup>, -NHCOR<sup>A</sup>, -NHSO<sub>2</sub>R<sup>A</sup>, and -NHR<sup>A</sup>, wherein R<sup>A</sup> is

(C<sub>1</sub>-C<sub>6</sub>)alkyl, cycloalkyl, phenyl, 2-, 3- or 4-pyridyl, N- or 2-, 3- or 4-piperidyl, N- or 2- or 3- piperazyl group; or

(ii) may be substituted by -NR<sup>A</sup>R<sup>B</sup>, -CONR<sup>A</sup>R<sup>B</sup> or -SO<sub>2</sub>NR<sup>A</sup>R<sup>B</sup>, wherein R<sup>A</sup> and R<sup>B</sup> are independently

(C<sub>1</sub>-C<sub>6</sub>)alkyl, cycloalkyl, phenyl, 2-, 3- or 4-pyridyl, N- or 2-, 3- or 4-piperidyl, N- or 2- or 3- piperazyl group,

or when taken together with the N atom to which they are attached R<sup>A</sup> and R<sup>B</sup> form a 5 to 7 membered aromatic or non- aromatic ring, which ring (a) may contain additional heteroatoms selected from N, O and S, and in which any S atom may be oxidised as a sulphonyl or sulfoxide, and (b) may be substituted on a ring carbon or heteroatom by one or more of the substituents listed under (i) above.

2. A method for the treatment of bacterial or protozoal infections in humans and non-human mammals, which comprises administering to a subject suffering such infection an antibacterially or antiprotozoally effective dose of a compound of formula (I) as defined in claim 1.

3. A compound of formula (I) as set forth in claim 1 wherein Z is a radical of formula  $-N(OH)CH(=O)$ , or a pharmaceutically or veterinarily acceptable salt, hydrate or solvate thereof.
  
4. A compound of formula (I) as set forth in claim 1 wherein Z is a radical of formula  $-C(=O)NH(OH)$  and  $R_4$  represents a radical  $R_5-(ALK)_p$ - wherein ALK is as defined in relation to  $R_2$ , p is 0 or 1, and  $R_5$  represents hydrogen, or a  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl, cycloalkyl, aryl, or heterocyclyl group which is substituted by  $-NR^A R^B$ ,  $-CONR^A R^B$  or  $-SO_2NR^A R^B$ , wherein  $R^A$  and  $R^B$  taken together with the N atom to which they are attached form a 5 to 7 membered aromatic or non-aromatic ring, which ring (a) may contain additional heteroatoms selected from N, O and S, and in which any S atom may be oxidised as a sulphonyl or sulfoxide, and (b) may be substituted by a group selected from  $(C_1$ - $C_6$ )alkyl, phenyl, benzyl,  $(C_1$ - $C_6$ )alkoxy, phenoxy, hydroxy, mercapto,  $(C_1$ - $C_6$ )alkylthio, amino, halo, trifluoromethyl, cyano, nitro, oxo,  $-COOH$ ,  $-SO_2H$ ,  $-CONH_2$ ,  $-SO_2NH_2$ ,  $-COR^A$ ,  $-SOR^A$ ,  $-SO_2R^A$ ,  $-SO_2R^A$ ,  $-COOR^A$ ,  $-CONHR^A$ ,  $-SO_2NHR^A$ ,  $-NHCOR^A$ ,  $-NHSO_2R^A$ , and  $-NHR^A$ , wherein  $R^A$  is  $C_1$ - $C_6$ )alkyl, cycloalkyl, phenyl, 2-, 3- or 4-pyridyl, N- or 2-, 3- or 4-piperidyl, N- or 2- or 3- piperazyl group, or a pharmaceutically or veterinarily acceptable salt, hydrate or solvate thereof.
  
5. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein  $R_1$  is hydrogen.
  
6. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein  $R_2$  is:  
 optionally substituted  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_6$  alkenyl,  $C_3$ - $C_6$  alkynyl or cycloalkyl;  
  
 phenyl( $C_1$ - $C_6$  alkyl)-, phenyl( $C_3$ - $C_6$  alkenyl)- or phenyl( $C_3$ - $C_6$  alkynyl)-  
 optionally substituted in the phenyl ring;

cycloalkyl(C<sub>1</sub>-C<sub>6</sub> alkyl)-, cycloalkyl(C<sub>3</sub>-C<sub>6</sub> alkenyl)- or cycloalkyl(C<sub>3</sub>-C<sub>6</sub> alkynyl)- optionally substituted in the cycloalkyl ring;

heterocyclyl(C<sub>1</sub>-C<sub>6</sub> alkyl)-, heterocyclyl(C<sub>3</sub>-C<sub>6</sub> alkenyl)- or heterocyclyl(C<sub>3</sub>-C<sub>6</sub> alkynyl)- optionally substituted in the heterocyclyl ring; or

CH<sub>3</sub>(CH<sub>2</sub>)<sub>p</sub>O(CH<sub>2</sub>)<sub>q</sub>- or CH<sub>3</sub>(CH<sub>2</sub>)<sub>p</sub>S(CH<sub>2</sub>)<sub>q</sub>-, wherein p is 0, 1, 2 or 3 and q is 1, 2 or 3.

7. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein R<sub>2</sub> is methyl, ethyl, n- or iso-propyl, n- and iso-butyl, n-pentyl, iso-pentyl, 3-methyl-but-1-yl, n-hexyl, n-heptyl, n-octyl, methylsulfanylethyl, ethylsulfanylmethyl, 2-methoxyethyl, 2-ethoxyethyl, 2-ethoxymethyl, 3-hydroxypropyl, allyl, 3-phenylprop-3-en-1-yl, prop-2-yn-1-yl, 3-phenylprop-2-yn-1-yl, 3-(2-chlorophenyl)prop-2-yn-1-yl, but-2-yn-1-yl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclopentylethyl, cyclopentylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexylpropyl, furan-2-ylmethyl, furan-3-methyl, tetrahydrofuran-2-ylmethyl, tetrahydrofuran-2-ylmethyl, piperidinylmethyl, phenylpropyl, 4-chlorophenylpropyl, 4-methylphenylpropyl, 4-methoxyphenylpropyl, benzyl, 4-chlorobenzyl, 4-methylbenzyl, or 4-methoxybenzyl.

8. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein R<sub>2</sub> is n-propyl, n-butyl, n-pentyl, or cyclopentylmethyl.

9. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein R<sub>3</sub> is:

the characterising group of a natural α amino acid, for example benzyl, or 4-methoxyphenylmethyl, in which any functional group may be protected, any amino group may be acylated and any carboxyl group present may be amidated; or

a group  $-[\text{Alk}]_n\text{R}_9$  where Alk is a  $(\text{C}_1-\text{C}_6)$ alkylene or  $(\text{C}_2-\text{C}_6)$ alkenylene group optionally interrupted by one or more  $-\text{O}-$ , or  $-\text{S}-$  atoms or  $-\text{N}(\text{R}_{12})-$  groups [where  $\text{R}_{12}$  is a hydrogen atom or a  $(\text{C}_1-\text{C}_6)$ alkyl group],  $n$  is 0 or 1, and  $\text{R}_9$  is hydrogen or an optionally substituted phenyl, aryl, heterocyclyl, cycloalkyl or cycloalkenyl group or (only when  $n$  is 1)  $\text{R}_9$  may additionally be hydroxy, mercapto,  $(\text{C}_1-\text{C}_6)$ alkylthio, amino, halo, trifluoromethyl, nitro,  $-\text{COOH}$ ,  $-\text{CONH}_2$ ,  $-\text{COOR}^A$ ,  $-\text{NHCOR}^A$ ,  $-\text{CONHR}^A$ ,  $-\text{NHR}^A$ ,  $-\text{NR}^A\text{R}^B$ , or  $-\text{CONR}^A\text{R}^B$  wherein  $\text{R}^A$  and  $\text{R}^B$  are independently a  $(\text{C}_1-\text{C}_6)$ alkyl group; or

a benzyl group substituted in the phenyl ring by a group of formula  $-\text{OCH}_2\text{COR}_8$  where  $\text{R}_8$  is hydroxyl, amino,  $(\text{C}_1-\text{C}_6)$ alkoxy, phenyl $(\text{C}_1-\text{C}_6)$ alkoxy,  $(\text{C}_1-\text{C}_6)$ alkylamino, di $((\text{C}_1-\text{C}_6)$ alkyl)amino, phenyl $(\text{C}_1-\text{C}_6)$ alkylamino; or

a heterocyclic $(\text{C}_1-\text{C}_6)$ alkyl group, either being unsubstituted or mono- or di-substituted in the heterocyclic ring with halo, nitro, carboxy,  $(\text{C}_1-\text{C}_6)$ alkoxy, cyano,  $(\text{C}_1-\text{C}_6)$ alkanoyl, trifluoromethyl  $(\text{C}_1-\text{C}_6)$ alkyl, hydroxy, formyl, amino,  $(\text{C}_1-\text{C}_6)$ alkylamino, di $(\text{C}_1-\text{C}_6)$ alkylamino, mercapto,  $(\text{C}_1-\text{C}_6)$ alkylthio, hydroxy $(\text{C}_1-\text{C}_6)$ alkyl, mercapto $(\text{C}_1-\text{C}_6)$ alkyl or  $(\text{C}_1-\text{C}_6)$ alkylphenylmethyl; or

a group  $-\text{CR}_a\text{R}_b\text{R}_c$  in which:

each of  $\text{R}_a$ ,  $\text{R}_b$  and  $\text{R}_c$  is independently hydrogen,  $(\text{C}_1-\text{C}_6)$ alkyl,  $(\text{C}_2-\text{C}_6)$ alkenyl,  $(\text{C}_2-\text{C}_6)$ alkynyl, phenyl $(\text{C}_1-\text{C}_6)$ alkyl,  $(\text{C}_3-\text{C}_6)$ cycloalkyl; or

$\text{R}_c$  is hydrogen and  $\text{R}_a$  and  $\text{R}_b$  are independently phenyl or heteroaryl such as pyridyl; or

$\text{R}_c$  is hydrogen,  $(\text{C}_1-\text{C}_6)$ alkyl,  $(\text{C}_2-\text{C}_6)$ alkenyl,  $(\text{C}_2-\text{C}_6)$ alkynyl, phenyl $(\text{C}_1-\text{C}_6)$ alkyl, or  $(\text{C}_3-\text{C}_6)$ cycloalkyl, and  $\text{R}_a$  and  $\text{R}_b$  together with the carbon atom to which they are attached form a 3 to 8 membered cycloalkyl or

a 5- to 6-membered heterocyclic ring; or

$R_a$ ,  $R_b$  and  $R_c$  together with the carbon atom to which they are attached form a tricyclic ring (for example adamantyl); or

$R_a$  and  $R_b$  are each independently  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl, phenyl $(C_1-C_6)$ alkyl, or a group as defined for  $R_c$  below other than hydrogen, or  $R_a$  and  $R_b$  together with the carbon atom to which they are attached form a cycloalkyl or heterocyclic ring, and  $R_c$  is hydrogen, -OH, -SH, halogen, -CN, -CO<sub>2</sub>H,  $(C_1-C_4)$ perfluoroalkyl, -CH<sub>2</sub>OH, -CO<sub>2</sub> $(C_1-C_6)$ alkyl, -O $(C_1-C_6)$ alkyl, -O $(C_2-C_6)$ alkenyl, -S $(C_1-C_6)$ alkyl, -SO $(C_1-C_6)$ alkyl, -SO<sub>2</sub> $(C_1-C_6)$ alkyl, -S $(C_2-C_6)$ alkenyl, -SO $(C_2-C_6)$ alkenyl, -SO<sub>2</sub> $(C_2-C_6)$ alkenyl or a group -Q-W wherein Q represents a bond or -O-, -S-, -SO- or -SO<sub>2</sub>- and W represents a phenyl, phenylalkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkylalkyl,  $(C_4-C_6)$ cycloalkenyl,  $(C_4-C_6)$ cycloalkenylalkyl, heteroaryl or heteroarylalkyl group, which group W may optionally be substituted by one or more substituents independently selected from, hydroxyl, halogen, -CN, -CO<sub>2</sub>H, -CO<sub>2</sub> $(C_1-C_6)$ alkyl, -CONH<sub>2</sub>, -CONH $(C_1-C_6)$ alkyl, -CONH $(C_1-C_6)$ alkyl)<sub>2</sub>, -CHO, -CH<sub>2</sub>OH,  $(C_1-C_4)$ perfluoroalkyl, -O $(C_1-C_6)$ alkyl, -S $(C_1-C_6)$ alkyl, -SO $(C_1-C_6)$ alkyl, -SO<sub>2</sub> $(C_1-C_6)$ alkyl, -NO<sub>2</sub>, -NH<sub>2</sub>, -NH $(C_1-C_6)$ alkyl, -N $((C_1-C_6)$ alkyl)<sub>2</sub>, -NHCO $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl,  $(C_3-C_6)$ cycloalkyl,  $(C_4-C_6)$ cycloalkenyl, phenyl or benzyl.

10. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein  $R_3$  is methyl, ethyl, benzyl, 4-chlorobenzyl, 4-hydroxybenzyl, phenyl, cyclohexyl, cyclohexylmethyl, pyridin-3-ylmethyl, tert-butoxymethyl, naphthylmethyl, iso-butyl, sec-butyl, tert-butyl, 1-benzylthio-1-methylethyl, 1-methylthio-1-methylethyl, 1-mercapto-1-methylethyl, 1-methoxy-1-methylethyl, 1-hydroxy-1-methylethyl, 1-fluoro-1-methylethyl, hydroxymethyl, 2-hydroxyethyl, 2-carboxyethyl, 2-methylcarbamoyl, 2-carbamoyl, or 4-



aminobutyl.

11. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein  $R_3$  is tert-butyl.

12. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein  $R_4$  is:

optionally substituted  $C_1$ - $C_8$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_8$  alkynyl or cycloalkyl;

optionally substituted phenyl, biphenyl or naphthyl;

optionally substituted piperidinyl, cyclohexyl, cyclopentyl, oxazolyl, thiazepinyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl, thienyl, furanyl, pyrrolyl, imidazolyl, benzofuranyl, benzothienyl, benzimidazolyl, thiazolyl, benzothiazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrazinyl, or triazinyl.

phenyl( $C_1$ - $C_6$  alkyl)-, phenyl( $C_2$ - $C_6$  alkenyl)- or phenyl( $C_2$ - $C_6$  alkynyl)- optionally substituted in the phenyl ring;


cycloalkyl( $C_1$ - $C_8$  alkyl)-, cycloalkyl( $C_2$ - $C_6$  alkenyl)- or cycloalkyl( $C_2$ - $C_6$  alkynyl)- optionally substituted in the cycloalkyl ring;

heterocyclyl( $C_1$ - $C_8$  alkyl)-, heterocyclyl( $C_2$ - $C_6$  alkenyl)- or heterocyclyl( $C_2$ - $C_6$  alkynyl)- optionally substituted in the heterocyclyl ring.

13. A use as claimed in claim 1, a method as claimed in claim 2, or compound as claimed in claim 3 or claim 4 wherein  $R_4$  is: methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 2-phenylcycloprop-1-yl, benzyl, biphenyl-2-yl, biphenyl-3-yl, biphenyl-4-yl, 2-methoxyphenyl, 3-methoxyphenyl, 4-

methoxyphenyl, 2,3-dimethoxyphenyl, 2,4-dimethoxyphenyl, 3,4-dimethoxyphenyl, 2-phenoxyphenyl, 3-phenoxyphenyl, 4-phenoxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2-iodophenyl, 3-iodophenyl, 4-iodophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 3,4-dimethyl, 2-aminophenyl, 3-aminophenyl, 4-aminophenyl, 2-acetamidophenyl, 3-acetamidophenyl, 4-acetamidophenyl, 2-N,N-dimethylaminophenyl, 3-N,N-dimethylaminophenyl, 4-N,N-dimethylaminophenyl, 2-methanesulfonamidophenyl, 3-methanesulfonamidophenyl, 4-methanesulfonamidophenyl, 2-hydroxyphenyl, 3-hydroxyphenyl, 4-hydroxyphenyl, 2,3-dihydroxyphenyl, 2,4-dihydroxyphenyl, 3,4-dihydroxyphenyl, 2-thiophenolyl, 3-thiophenolyl, 4-thiophenolyl, 2-thioanisoyl, 3-thioanisoyl, 4-thioanisoyl, 1-naphthyl, 2-naphthyl, furan-2-yl, thien-2-yl, pyrrol-2-yl, 1-methylpyrrol-2-yl, imidazol-2-yl, 1-methylimidazol-2-yl, thiazol-2-yl, 5-phenylpyrrol-2-yl, 5-phenylfuran-2-yl, 5-phenylthien-2-yl, benzothiazol-2-yl, 1,2,4-oxadiazol-5-yl, 3-methyl-1,2,4-oxadiazol-5-yl, 3-phenyl-1,2,4-oxadiazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,2,4-thiadiazol-5-yl, 1,3,4-thiadiazol-2-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, N-oxides of pyridin-2-yl pyridin-3-yl and pyridin-4-yl, indol-2-yl, indol-3-yl, 1-methylindol-2-yl, 1-methylindol-3-yl, benzimidazol-2-yl, 1-methylbenzimidazol-2-yl, pyrazin-2-yl, 1,2-pyridazin-3-yl, 1,3-pyrimidin-2-yl, benzo[b]thien-2-yl, benzo[b]thien-3-yl, benzo[b]furan-2-yl, benzo[b]furan-3-yl, isoxazol-5-yl, quinolin-2-yl, quinolin-3-yl, isoquinolin-2-yl, isoquinolin-3-yl, 2-oxo-2-phenylethyl, diphenylmethyl, 4-N-methylaminophenyl, 4-N,N-dimethylcarboxamidophenyl, and 4-carboxyphenyl.

14. A use as claimed in claim 1, a method as claimed in claim 2, a compound as claimed in claim 3 or claim 4 wherein  $R_4$  is a phenyl group which is substituted by one of the following:



15. A compound which is specifically named and characterised in any of Examples 1 to 32 herein.

16. A pharmaceutical composition comprising a compound as claimed in claim 3 or claim 4, or any of claims 5 to 15 together with a pharmaceutically acceptable carrier.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/01337

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/165 C07C239/18 A61K31/395 A61K31/335 A61P31/04  
 C07D295/10 C07D233/64 C07D213/50 C07D307/81 C07C317/40  
 C07D307/52 C07D207/32 C07C311/08 C07D239/42 C07D239/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, CHEM ABS Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages   | Relevant to claim No. |
|----------|--|-----------------------|
| P,X      | WO 99 39704 A (BRITISH BIOTECH PHARMACEUTICALS)<br>12 August 1999 (1999-08-12)<br>cited in the application<br>page 31, line 5 - line 17<br>---   | 15, 16                |
| A        | JOHN P. DEVLIN ET AL.: "Studies Concerning the Antibiotic Actinonin. Part III. Synthesis of Structural Analogues of Actinonin by the Anhydride-Imide Method" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS I, 1975, pages 830-841, XP002141667 CHEMICAL SOCIETY, LETCHWORTH., GB page 830; table 4<br>---<br>-/-- | 1                     |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 July 2000

Date of mailing of the international search report

25/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Zervas, B

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 00/01337

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|----------|---|-----------------------|
| A        | FR 2 353 289 A (INSTITUT PASTEUR DE LYON)<br>30 December 1977 (1977-12-30)<br>claims; examples<br>---                                     | 1                     |
| A        | US 3 166 587 A (JACK BERNSTEIN ET AL.)<br>19 January 1965 (1965-01-19)<br>claims; examples<br>---   | 1                     |
| A        | WO 97 03783 A (BRITISH BIOTECH<br>PHARMACEUTICALS)<br>6 February 1997 (1997-02-06)<br>cited in the application<br>claims; examples<br>--- | 1,15                  |
| A        | WO 95 33709 A (F. HOFFMANN-LA ROCHE)<br>14 December 1995 (1995-12-14)<br>cited in the application<br>claims; examples<br>-----            | 1,15                  |

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1, 2, 5-14 (all in part)

Present claims 1, 2 and 5-14 relate to the use of an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds claimed in claims 3, 4, 15 and 16 and their use.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/GB 00/01337

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---------------------|----------------------------|---------------------|
| WO 9939704 A                              | 12-08-1999          | AU 2529299 A               | 23-08-1999          |
| FR 2353289 A                              | 30-12-1977          | NONE                       |                     |
| US 3166587 A                              | 19-01-1965          | NONE                       |                     |
| WO 9703783 A                              | 06-02-1997          | AU 6663396 A               | 18-02-1997          |
|   |                     | EP 0865339 A               | 23-09-1998          |
|   |                     | GB 2318353 A,B             | 22-04-1998          |
|   |                     | JP 11510140 T              | 07-09-1999          |
| WO 9533709 A                              | 14-12-1995          | AU 2735995 A               | 04-01-1996          |
|   |                     | EP 0763013 A               | 19-03-1997          |
|   |                     | JP 10500965 T              | 27-01-1998          |
|   |                     | US 6066662 A               | 23-05-2000          |